

CASE REPORT

Left bundle branch block, atrioventricular block, torsade de pointes and long QT syndrome: Is this too much for a rare cardiomyopathy?☆

Bruno Rodrigues*, Emanuel Correia, Luís Ferreira Santos, Davide Moreira, Anne Delgado, Pedro Gama, António Costa, João Pipa, Oliveira Santos

Serviço de Cardiologia, Hospital São Teotónio, Centro Hospitalar Tondela-Viseu, Viseu, Portugal

Received 3 October 2011; accepted 4 June 2012

Available online 6 June 2013

KEYWORDS

Left ventricular noncompaction;
Atrioventricular block;
Torsade de pointes

Abstract Left ventricular noncompaction (LVNC) is now recognized as a distinct form of cardiomyopathy with a clinical presentation and natural history of its own. Common manifestations of LVNC include heart failure, ventricular arrhythmias and embolic events, but serious atrioventricular conduction disturbances are rarely reported in the literature. The authors describe the case of a 40-year-old woman who went to the emergency department due to syncope. The ECG revealed left bundle branch block (LBBB) and 2:1 atrioventricular block (AVB) and the patient was admitted for pacemaker implantation. During hospitalization she developed torsade de pointes and complete AVB with increased QTc. The echocardiogram showed images compatible with LVNC. This case provides additional evidence that LVNC may be complicated by 2:1 (or complete) AVB, intraventricular conduction disturbances (LBBB) and repolarization abnormalities (long QT). This combination of electrocardiographic changes has rarely been reported in the literature. We describe a series of affected patients, focusing on electrocardiographic characteristics.

© 2011 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L. All rights reserved.

PALAVRAS-CHAVE

Ventrículo esquerdo não compactado;
Bloqueio auriculoventricular;
Torsade de pointes

Bloqueio completo de ramo esquerdo, bloqueio auriculoventricular, torsade de pointes e QT longo: será demasiado para uma rara miocardiopatia?

Resumo O ventrículo esquerdo não compactado (VENC) tem sido reconhecido como uma forma distinta de miocardiopatia com uma apresentação clínica e história natural próprias. As manifestações comuns de VENC são insuficiência cardíaca, arritmias ventriculares e eventos embólicos. A presença de perturbações da condução auriculoventricular de maior gravidade encontra-se raramente descrita na literatura. Os autores descrevem o caso clínico de uma

☆ Please cite this article as: Rodrigues B, et al. Bloqueio completo de ramo esquerdo, bloqueio auriculoventricular, torsade de pointes e QT longo: será demasiado para uma rara miocardiopatia? Rev Port Cardiol. 2013. <http://dx.doi.org/10.1016/j.repc.2012.06.018>.

* Corresponding author.

E-mail address: Onurb80@sapo.pt (B. Rodrigues).

paciente com 40 anos de idade, que recorreu ao Serviço de Urgência por síncope. O ECG revelou bloqueio completo de ramo esquerdo do feixe de His e bloqueio auriculoventricular (BAV) 2:1, tendo sido internada para implantação de *pacemaker*. Durante o internamento, desenvolve fenómenos de *torsade de pointes* e de BAV completo com aumento do intervalo QTc. O ecocardiograma revelou imagens compatíveis com VENC. Com este caso, os autores fornecem evidência adicional de que a presença de VENC pode ser complicada por BAV 2:1 (ou completo), perturbações da condução intraventricular (BCRE) e da repolarização (QT longo), alterações eletrocardiográficas concomitantes pouco descritas na literatura. Descreve-se uma série de pacientes afetados, com foco nas características eletrocardiográficas.

© 2011 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L. Todos os direitos reservados.

Introduction

Left ventricular noncompaction (LVNC) is a rare cardiomyopathy originally described in children^{1,2} but whose phenotypic characteristics are increasingly documented in adults.³⁻⁷ The most common symptoms of LVNC are heart failure (HF), ventricular arrhythmias and embolic events, but atrioventricular conduction disturbances are rarely reported in the literature.⁸⁻¹⁰

Case report

A 40-year-old woman went to the emergency department due to syncope during a meal, without prodrome and with spontaneous recovery, associated with progressive fatigue for around a month. She reported no chest pain or dyspnea. Personal history included complete left bundle branch block (LBBB) detected on a routine electrocardiogram (ECG) as part of a medical check-up at work three months before. There was no family history of sudden death and she was not taking any medication.

Physical examination showed the patient to be eupneic with no signs of respiratory difficulty. Cardiac auscultation revealed rhythmic sounds and no murmurs but bradycardia (40 bpm); pulmonary auscultation detected no abnormalities. Blood pressure was 100/75 mmHg. The rest of the physical examination was normal.

Laboratory tests, including blood and biochemical analysis, showed no relevant alterations and biomarkers of myocardial necrosis were normal.

A 12-lead ECG revealed second-degree 2:1 atrioventricular block (AVB) with ventricular response of ± 45 bpm and LBBB, and the patient was accordingly admitted for pacemaker implantation (Figure 1).

During hospitalization she developed cardiac arrest due to torsade de pointes (TdP) (Figure 2), reverted by cardiac massage. The ECG revealed 2:1 AVB (with a period of complete AVB), heart rate of ± 45 bpm and increased QTc interval (Figure 3).

Echocardiography performed in the context of monitoring in the cardiac care unit showed a normal-size left ventricle with mildly impaired global systolic function (ejection fraction $\pm 45\%$) and images compatible with LVNC at the

apex (pronounced trabeculations and ratio of trabeculated to non-trabeculated layers >2 , with blood flow on color Doppler) (Figures 4 and 5). Coronary angiography revealed no coronary disease.

A dual-chamber cardioverter-defibrillator was accordingly implanted for secondary prevention of sudden death.

The patient was discharged clinically stable and is currently well with no signs of HF. She is pacing dependent but no shocks have been detected. Screening of first-degree relatives showed normal ECG but echocardiography in her children showed images compatible with inferior and lateral LVNC (Figure 6), although high-resolution ECG showed no abnormalities. The results of the patient's genetic study are awaited.

Discussion

Patients with LVNC usually have impaired left ventricular systolic function and generally present with symptoms of HF.^{1,2,4} Other associated cardiac alterations include endocardial thrombi, leading to systemic embolism, and arrhythmias. The high incidence of arrhythmias in LVNC is not explained by a known morphological abnormality, but it has been suggested that myocardial ischemia may play an important role in the development of systolic dysfunction and arrhythmias in these patients, as is seen in ischemic heart disease.¹¹ A variety of rhythm disturbances have been reported in LVNC, of which ventricular arrhythmias are the most common; they may be refractory to medical treatment and can be fatal.^{1-4,11-15}

The literature shows that 88–94% of LVNC patients have alterations on the baseline ECG.^{2,3,16} In a series by Oechslin et al.,⁴ 94% of the 34 adults studied presented abnormal ECGs and 56% had intraventricular conduction defects, most of them identified in patients who had died. In a series by Steffel et al.¹⁶ (n=78), only 13% of the patients had a normal ECG, while two (3%) had complete AVB block, 19% had LBBB, and 50% had prolonged QTc.

Wolff–Parkinson–White syndrome (WPW) and associated tachycardias have also been described in association with LVNC.^{1,2,14,17} Ichida et al.² reported WPW syndrome in four of 27 patients with LVNC, and explained this association as a failed regression of developmental embryologic atrioventricular anatomical and electrical continuity during




Figure 1 ECG showing second-degree 2:1 atrioventricular block with left bundle branch block.

embryonic development in the non-compacted myocardium. Supraventricular tachycardias associated with WPW syndrome can thus trigger fatal ventricular tachycardia and fibrillation in patients with LVNC.^{1,17}

In our series⁶ (n=20) (Table 1), there was one case of 2:1/intermittent complete AVB, three cases of LBBB, and six cases of increased QTc interval; two cases presented as ventricular tachycardia. Only two patients (10%) had a normal ECG, a similar proportion to that reported by Oechslin et al. There were no cases of WPW syndrome.

The cause of intraventricular conduction disturbances and repolarization abnormalities in LVNC is unknown. Ichida et al.² suggest that conduction disturbances such as LBBB and repolarization abnormalities such as increased QTc result from progressive endocardial fibrosis, and are often observed in adult patients because fibrotic changes develop gradually.⁸

With regard to the atrioventricular conduction disturbances observed in this case (second-degree/intermittent

complete AVB), it is suspected that they may be caused by histological alterations in the conduction system itself⁸ alongside the endomyocardial fibrotic process. However, long QT complicating AVB, with concomitant ventricular arrhythmias such as TdP, is also frequently seen in association with conduction disturbances of any etiology, and may be aggravated by the presence of LVNC for the reasons described above.

The probability of patients under antiarrhythmic medication and/or with ionic imbalance developing long QT and TdP is increased by bradycardia due to reverse use-dependence of potassium channels.¹⁸ However, the role of bradycardia itself in the genesis of TdP is less clear.^{19,20} This question was addressed in a study¹⁹ of 14 patients with complete AVB, six of whom had a history of TdP. There was no difference in the frequency of ventricular escape rhythm between the two groups, but QTc was significantly longer in those with a history of TdP (0.59 vs. 0.48 s). After pacemaker implantation, QTc was still longer in those with a history of TdP,




Figure 2 Torsade de pointes.




Figure 3 2:1 atrioventricular block (heart rate \pm 45 bpm) with prolonged QTc interval (\pm 690 ms).

Table 1 ECG abnormalities in our series of patients with LVNC (n=20; mean age 52.8 years; 71% male).

	n	Percentage
<i>Rhythm</i>		
Sinus	15	75%
Atrial fibrillation	5	25%
<i>Ventricular arrhythmia at admission</i>		
No	18	90%
Yes	2	10%
<i>AV conduction disturbances</i>		
No	15	75%
1st-degree AVB	4	20%
2nd-degree/complete AVB	1	5%
<i>Intraventricular conduction disturbances</i>		
No	13	65%
Left anterior hemiblock	3	15%
RBBB	1	5%
LBBB	3	1%
<i>Repolarization disturbances – stress^a</i>		
No	10	50%
Yes	7	35%
<i>Repolarization disturbances – ischemia^a</i>		
No	13	65%
Yes	4	20%
<i>Long QTc interval</i>		
No	14	70%
Yes	6	30%
<i>Signs of left ventricular hypertrophy^a</i>		
No	14	70%
Yes	3	15%

AV: atrioventricular; AVB: atrioventricular block; LBBB: left bundle branch block; RBBB: incomplete right bundle branch block.

^a Excluding patients with LBBB.



Figure 4 Echocardiogram: parasternal short-axis view showing images compatible with left ventricular non-compaction at the apex.




Figure 5 Echocardiogram. Left: apical 4-chamber view showing images compatible with left ventricular non-compaction at the apex; right: three-dimensional echocardiogram showing detail of trabeculations (arrow).




Figure 6 (A and B) Echocardiograms of the patient's children showing images compatible with left ventricular non-compaction.

irrespective of whether the lower rate limit was set at 60 bpm (0.55 vs. 0.50 s) or 50 bpm (0.70 vs. 0.53 s).

Conclusion

LVNC is often difficult to diagnose, but a high degree of clinical awareness is necessary due to its poor prognosis.

To the best of our knowledge, this is the first reported case of LVNC associated with LBBB, second-degree/intermittent complete AVB, ventricular arrhythmia and long QT syndrome. We therefore propose that a diagnosis of LVNC be considered in relatively young individuals who present with high-degree AVB and LBBB and with a relevant clinical history.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data and that all the patients included in the

study received sufficient information and gave their written informed consent to participate in the study.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Chin TK, Preloff JK, Williams RG, et al. Isolated noncompaction of left ventricular myocardium. A study of eight cases. *Circulation*. 1990;82:507–13.
- Ichida F, Hamamichi Y, Miyawaki T, et al. Clinical features of isolated noncompaction of the ventricular myocardium: long-term clinical course, hemodynamic properties, and genetic background. *J Am Coll Cardiol*. 1999;34:233–40.
- Ritter M, Oechslin E, Sütsch G, et al. Isolated noncompaction of the myocardium in adults. *Mayo Clin Proc*. 1997;72:26–31.
- Oechslin EN, Attenhofer Jost CH, Rojas JR, et al. Long-term follow-up of 34 adults with isolated left ventricular noncompaction: a distinct cardiomyopathy with poor prognosis. *J Am Coll Cardiol*. 2000;36:493–500.
- Daimon Y, Watanabe S, Takeda S, et al. Two-layered appearance of noncompaction of the ventricular myocardium on magnetic resonance imaging. *Circ J*. 2002;66:619–21.
- Correia E, Rodrigues B, Santos L, et al. Miocárdio não-compactado isolado: caracterização e seguimento de população afectada. *Rev Port Cardiol*. 2011;30:323–31.
- Jenni R, Oechslin E, Schneider J, et al. Echocardiographic and pathoanatomical characteristics of isolated left ventricular non-compaction: a step towards classification as a distinct cardiomyopathy. *Heart*. 2001;86:666–71.
- Taniguchi M, Hioka T, Maekawa K, et al. Adult case of isolated ventricular noncompaction discovered by complete atrioventricular block. *Circ J*. 2004;68:873–5.
- Yildiz A, Ozeke O, Akyol S, et al. Biventricular myocardial non-compaction presenting with complete atrioventricular block. *Int J Cardiol*. 2009;132:e34–6.
- Nascimento BR, Vidigal DF, Carneiro RD, et al. Complete atrioventricular block as the first manifestation of noncompaction of the ventricular myocardium. *Pacing Clin Electrophysiol*. 2011. <http://dx.doi.org/10.1111/j.1540-8159.2011.03054.x>.
- Celiker A, Ozkutlu S, Dilber E, et al. Rhythm abnormalities in children with isolated ventricular noncompaction. *Pacing Clin Electrophysiol*. 2005;28:1198–202.
- Robida A, Hajar HA. Ventricular conduction defect in isolated noncompaction of the ventricular myocardium. *Pediatr Cardiol*. 1996;17:189–91.
- Ozkutlu S, Ayabakan C, Celiker A, et al. Noncompaction of ventricular myocardium: a study of twelve patients. *J Am Soc Echocardiogr*. 2002;15:1523–8.
- Elsershari H, Okutan V, Celiker A. Isolated noncompaction of ventricular myocardium. *Cardiol Young*. 2001;11:472–5.
- Wald R, Veldtman G, Golding F, et al. Determinants of outcome in isolated ventricular noncompaction in childhood. *Am J Cardiol*. 2004;94:1581–4.
- Steffel J, Kobza R, Oechslin E, et al. Electrocardiographic characteristics at initial diagnosis in patients with isolated left ventricular noncompaction. *Am J Cardiol*. 2009;104:984–9.
- Yasukawa K, Terai M, Honda A, et al. Noncompaction of ventricular myocardium associated with fatal ventricular fibrillation. *Pediatr Cardiol*. 2001;22:512–4.

18. Yang T, Roden DM. Extracellular potassium modulation of drug block of IKr. Implications for torsade de pointes and reverse use-dependence. *Circulation*. 1996;93:407-11.
19. Kurita T, Ohe T, Marui N, et al. Bradycardia-induced abnormal QT prolongation in patients with complete atrioventricular block with torsades de pointes. *Am J Cardiol*. 1992;69:628-33.
20. Furushima H, Niwano S, Chinushi M, et al. Relation between bradycardia dependent long QT syndrome and QT prolongation by disopyramide in humans. *Heart*. 1998;79:56-8.