



LETTER TO THE EDITOR

Reply to the letter “The relation between inflammation and coronary artery ectasia”



Resposta à carta «A relação entre a inflamação e a ectasia arterial coronária»

Dear Editor,

We would like to thank Balta et al.¹ for their interest in our article.² In our study, we evaluated the frequency of major cardiovascular risk factors and serum levels of gamma-glutamyltransferase (GGT) and high-sensitivity C-reactive protein (hs-CRP) in a relatively large population of patients with isolated coronary artery ectasia (CAE). We found that CAE can be independently and positively associated with obesity, GGT and hs-CRP levels, but inversely with diabetes. Moreover, its severity may be related to GGT and hs-CRP levels.²

Coronary artery ectasia is usually defined as dilation exceeding 1.5 times the diameter of adjacent normal segments in epicardial coronary arteries.^{3,4} Its underlying causes are poorly understood, but it has frequently been considered as a variant of atherosclerotic vascular disease.³ Atherosclerosis is regarded as a low-grade inflammatory process⁵ and thus it is not surprising that there may be a close association between inflammation and CAE. As reported by Balta et al.,¹ various studies have shown such an association.^{6–10} Moreover, inflammation may be related to severity of CAE. However, the extent of CAE can be differently defined. Markis et al. categorized CAE in four types, with decreasing severity of CAE from type I to IV according to its topographical extent in the major coronary arteries.³ Based on this definition, types I, II and III CAE can be considered severe ectasia. By contrast, we defined severe CAE as diffuse involvement (≥ 2 segments) in at least two vessels. This definition corresponds to type I CAE according to Markis’s classification. Our definition has also been used in previous studies.^{8,9}

We agree with Balta et al. that some disorders such as obesity, thyroid dysfunction and malignancy can trigger an

inflammatory state in which inflammatory markers like hs-CRP may be elevated. We excluded patients with thyroid dysfunction and known malignancies and inflammatory diseases from the study. With obesity, we consecutively included CAE patients in the study and selected age- and gender-matched controls with normal coronary arteries. Although mean body mass index was comparable between CAE patients and controls, the obesity rate was slightly higher in CAE patients (16% vs. 9%, $p=0.06$) and obesity was positively associated with CAE. We think that inflammatory activity may contribute to the development of CAE in obese patients.

Serum GGT, a major antioxidant, can oxidize low-density lipoprotein cholesterol, and has a role in the pathogenesis of atherosclerosis.¹¹ GGT can also act as a proinflammatory protein in atherogenesis and is associated with atherosclerotic risk factors including obesity, dyslipidemia, metabolic syndrome, hypertension and diabetes.^{12,13} Primary hepatic diseases such as alcoholic or viral hepatitis and Gilbert syndrome can lead to elevated transaminases, GGT, and bilirubin levels in hepatic function tests. We did not enroll patients who had previously known hepatic disease and who regularly drank ethyl alcohol. Accordingly, we think that the relatively elevated levels of GGT may be associated with CAE itself rather than other disorders or situations.

In conclusion, there is growing evidence that an inflammatory state may be activated in CAE patients and that its markers may be elevated in CAE patients as in obstructive coronary artery disease. As an oxidative stress and proinflammatory marker, GGT may also have a role in the pathogenesis of CAE. This consideration needs future studies.

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Balta S, Demir M, Ozturk C, et al. The relation between inflammation and coronary artery ectasia. *Rev Port Cardiol.* 2016;35:553–4.
- Dogan A, Arslan A, Yucel H, et al. Gamma glutamyltransferase, inflammation and cardiovascular risk factors in isolated coronary artery ectasia. *Rev Port Cardiol.* 2016;35:33–9.

DOI of original article:
<http://dx.doi.org/10.1016/j.repc.2016.04.011>

<http://dx.doi.org/10.1016/j.repc.2016.08.002>

0870-2551/© 2016 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. All rights reserved.

3. Markis J, Joffe C, Cohn P, et al. Clinical significance of coronary artery ectasia. *Am J Cardiol.* 1976;37:217–22.
4. Swanton RH, Thomas ML, Coltart DJ, et al. Coronary artery ectasia – a variant of occlusive coronary arteriosclerosis. *Br Heart J.* 1978;40:393–400.
5. Libby P. Inflammation in atherosclerosis. *Nature.* 2002;420:868–74.
6. Tokgozoglul L, Ergene O, Kinay O, et al. Plasma interleukin-6 levels are increased in coronary artery ectasia. *Acta Cardiol.* 2004;59:515–9.
7. Turhan H, Erbay AR, Yasar AS, et al. Comparison of C-reactive protein levels in patients with coronary artery ectasia versus patients with obstructive coronary artery disease. *Am J Cardiol.* 2004;94:1303–6.
8. Finkelstein A, Michowitz Y, Abashidze A, et al. Temporal association between circulating proteolytic, inflammatory and neurohormonal markers in patients with coronary ectasia. *Atherosclerosis.* 2005;179:353–9.
9. Dogan A, Tuzun N, Turker Y, et al. Matrix metalloproteinases and inflammatory markers in coronary artery ectasia: their relationship to severity of coronary artery ectasia. *Coron Artery Dis.* 2008;19:559–63.
10. Li JJ, Nie SP, Qian XW, et al. Chronic inflammatory status in patients with coronary artery ectasia. *Cytokine.* 2009;46: 61–4.
11. Emdin M, Pompella A, Paolicchi A. Gamma glutamyltransferase, atherosclerosis, and cardiovascular disease: triggering oxidative stress within the plaque. *Circulation.* 2005;112:2078–80.
12. Lee DH, Jacobs DR, Gross M, et al. Gamma-glutamyltransferase is a predictor of incident diabetes and hypertension: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Clin Chem.* 2003;49:1358–66.
13. Lee DS, Evans JC, Robins SJ, et al. Gamma glutamyltransferase and metabolic syndrome, cardiovascular disease, and mortality risk: the Framingham Heart Study. *Arterioscler Thromb Vasc Biol.* 2007;27:127–33.

Abdullah Dogan, Akif Arslan*

Cardiology Department, Medical School, Suleyman Demirel University, Isparta, Turkey

* Corresponding author.

E-mail address: dr.akifarslan@hotmail.com (A. Arslan).