

IMAGE IN CARDIOLOGY

Acute L waves in heart failure

Ondas Lambda agudas na insuficiência cardíaca

Juan Diego Sánchez Vega*, Sergio García Gómez, Manuel García de Yébenes Castro

Clínica Universidad de Navarra, Madrid, Spain

Received 29 May 2023; accepted 5 June 2023

Available online 2 November 2023



L waves are an unusual finding in the analysis of mitral valve pulsed-wave Doppler. This wave is recorded as flow between the left atrium and left ventricle during diastole, due to elevated left ventricular (LV) filling pressures.^{1–3} In this situation, triphasic mitral filling can be observed, with the presence of mid-diastolic L waves.

We present the case of a 71-year-old female with preserved LV ejection fraction (LVEF) and moderate ventricular hypertrophy secondary to hypertensive cardiomyopathy, according to a cardiac magnetic resonance study performed the year before. The patient had an episode of acute pulmonary edema during an admission for malaria, with complications of fever, acute kidney disease and hypertensive crisis. During the initial assessment, a transthoracic echocardiogram was performed, which showed high LV filling pressure (E/E' ratio 23), systolic pulmonary pressure of 60 mmHg estimated by tricuspid regurgitation Doppler, and L

waves on mitral pulsed-wave Doppler (Figure 1). Laboratory tests showed NT-proBNP of 1962 pg/ml.

After treatment with antihypertensive drugs and diuretics, following diuresis of 12 l in three days, the patient had normal LV filling pressure (E/E' ratio 12) and systolic pulmonary pressure of 32 mmHg, and the L wave had disappeared on the echocardiogram, which showed an impaired relaxation pattern in LV filling (E<A wave). NT-proBNP had normalized (19 pg/ml). We also observed an improvement in left atrial contractility, measured by left atrial conduit strain (initial –8.4%, after treatment –12%) (Videos 1 and 2), and a decrease in left atrial volume (from 52.1 to 31.7 ml).

The patient was discharged in good condition after recovery of renal function and adjustment of antihypertensive therapy.

In this paper, we report the presence of L waves during an episode of acute pulmonary edema in a patient with

* Corresponding author.

E-mail address: jdsanchez@unav.es (J.D. Sánchez Vega).




Figure 1 (A and B) Chest radiography during the episode of heart failure after treatment; (C) L wave between rapid left ventricular (LV) filling (E wave) and atrial contraction (A wave); (D) no L wave after decongestion, with impaired LV relaxation pattern (E<A).

preserved LVEF, in the context of high LV filling pressure. Recognizing this pattern of diastolic filling can be helpful in the rapid diagnosis and treatment of similar cases of heart failure.

Conflicts of interest

The authors have no conflicts of interest to declare.

Appendix A. Supplementary data

Supplementary material associated with this article can be found in the online version at doi:10.1016/j.rep.2023.06.010.

References

1. Kumar V, Jose J, Jose VJ. L wave in echo Doppler. Indian Heart J. 2014;66:392–3.
2. Lam CSP, Han L, Ha JW, et al. The mitral L wave: a marker of pseudonormal filling and predictor of heart failure in patients with left ventricular hypertrophy. J Am Soc Echocardiogr. 2005;18:336–41.
3. Ha JW, Ahn JA, Moon JY, et al. Triphasic mitral inflow velocity with mid-diastolic flow: the presence of mid-diastolic mitral annular velocity indicates advanced diastolic dysfunction. Eur J Echocardiogr. 2006;7:16–21.