

# Journal Pre-proof

Navigating the challenges of ST-segment elevation myocardial infarction in women: A closer look

Marisa Trabulo Ana Teresa Timóteo



PII: S0870-2551(25)00031-9

DOI: <https://doi.org/doi:10.1016/j.repc.2025.01.002>

Reference: REPC 2406

To appear in: *Revista Portuguesa de Cardiologia*

Received Date: 10 January 2025

Accepted Date: 20 January 2025

Please cite this article as: Trabulo M, Timóteo AT, Navigating the challenges of ST-segment elevation myocardial infarction in women: A closer look, *Revista Portuguesa de Cardiologia* (2025), doi: <https://doi.org/10.1016/j.repc.2025.01.002>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier España, S.L.U. on behalf of Sociedade Portuguesa de Cardiologia.

## Navigating the challenges of ST-segment elevation myocardial infarction in women: A closer look

### Dificuldades no enfarte com elevação do segmento ST nas mulheres: um olhar atento

Marisa Trabulo<sup>1,2,3\*</sup>, Ana Teresa Timóteo<sup>2,4\*</sup>

1 – Serviço Cardiologia, Hospital Santa Cruz, Unidade Local Saúde Lisboa Ocidental, Carnaxide, Portugal

2 – NOVA Medical School, Universidade Nova de Lisboa, Lisboa, Portugal

3 - Hospital da Luz, Lisboa, Portugal

4 – Serviço de Cardiologia, Hospital Santa Marta, Unidade Local Saúde São José, Lisboa, Portugal

\*Corresponding author:

E-mail address: marisatrabulo@hotmail.com (M. Trabulo)

**\*These authors contributed equally to this work and share the first authorship.**

Cardiovascular diseases are the leading cause of death in Europe and in developed countries, particularly in women<sup>1</sup>. There are several factors contributing to a worst prognosis in women, but one of the most important is the fact that women, family members and clinicians tend to under-recognize clinical symptoms and signs of heart disease in female patients<sup>2</sup>. This will cause underdiagnosis, significant delays in treatment and frequently, undertreatment<sup>2</sup>. Moreover, women are usually under-represented in clinical trials, and evidence is less strong in women<sup>2</sup>. Notably, in women, classical risk factors have a higher impact on outcome compared to men. In addition, there are other risk factors that are more frequent in women, with some that are exclusive to women (due to their specific biology) that contribute to this heightened female risk<sup>2</sup>.

In the context of acute coronary syndromes, there are significant disparities between men and women. These differences refer to anatomical, physiological, biological, and psychosocial factors which affect each other<sup>3</sup>. Situations of myocardial infarction with non-obstructive coronary arteries (MINOCA), due to microcirculation abnormalities, coronary spasm, spontaneous coronary dissections or even Takotsubo syndrome are more frequently found in women<sup>3,4</sup>.

A paper published in 2020 with data from the Portuguese Registry of Acute Coronary Syndromes (ProACS) showed very elegantly that in Portugal, as in other countries, obesity, hypertension and diabetes are more prevalent in

women and that they have more frequently non-ST segment elevation myocardial infarction, with atypical symptoms<sup>5</sup>. Delays in treatment are more frequent, with more complications, such as major bleeding, heart failure, atrial fibrillation, mechanical complications, cardiogenic shock, stroke, and ultimately, higher in-hospital mortality<sup>5</sup>. They also received less guideline recommended secondary prevention therapies<sup>5</sup>. Therefore, the landscape in Portugal regarding ACS seems to mirror what has been described in other European countries.

In the study published in the current issue of the Portuguese Journal of Cardiology, Gonçalves et al conducted a retrospective analysis of a large cohort of ST-elevation myocardial infarction (STEMI) patients included in the ProACS registry, between October 2010 and 2022<sup>6</sup>. The co-primary endpoints were in-hospital and one-year mortality. In this analysis, it was the author's option to include only STEMI patients, to homogenize the population studied, reducing potential confounding factors related to population heterogeneity.

Women were significantly under-represented (25.7%), and this is a very significant finding, considering that the data were obtained from a registry and not from a clinical trial. Significant sex-related differences in Portuguese STEMI patients were found. Compared to men, women were older and had a worse cardiovascular risk profile. They were less likely to receive reperfusion therapy as well as other medication with an impact on prognosis. In-hospital mortality was higher in women.

In line with other publications, but specifically addressing the Portuguese population, this study confirms that there is still a gap in the diagnosis and treatment of acute coronary syndromes in women, and specifically in STEMI patients, in whom fewer gender-differences would be expected, since the diagnosis and management is generally more straightforward than in the general population with ACS.

Every effort should be made toward achieving an equal and high standard of care. Education campaigns targeting the general population and healthcare professionals may help reduce this sex bias. Noncardiac comorbidities should also be strongly accounted for.

## References

- 1 - Timmis A, Vardas P, Townsend N, et al. European Society of Cardiology: cardiovascular disease statistics 2021. *Eur Heart J* 2022;43: 716–99.
- 2 - Timóteo AT, Ribeiros R, Calé R, et al. Cardiovascular diseases in women - Consensus document of the Sociedade Portuguesa de Cardiologia, Sociedade Portuguesa de Ginecologia, Sociedade Portuguesa de Obstetrícia e Medicina Materno-Fetal, Sociedade Portuguesa de Contraceção e Associação Portuguesa de Medicina Geral e Familiar. *Rev Port Cardiol* 2023; 42: 1001-1015.
- 3 – Almeida A, Grapsa J, Gimelli A, et al. Cardiovascular multimodality imaging in women: a scientific statement of the European Association of Cardiovascular Imaging of the European Society of Cardiology. *Eur Heart J Cardiovasc Imag* 2024; 25: doi:/10.1093/ehjci/jeae013
- 4 - Brízido C, Madeira S, Silva C, et al. Spontaneous coronary artery dissection: A review for clinical and interventional cardiologists. *Rev Port Cardiol* 2023; 42: 269-276. doi: 10.1016/j.repc.2022.03.008.
- 5 - Roque D, Ferreira J, Monteiro S, et al. Understanding a woman's heart: Lessons from 14 177 women with acute coronary syndrome. *Rev Port Cardiol* 2020; 39: 57-72.
- 6 – C. Miguel Gonçalves, M. Carvalho, A. Vazão et al., Sex-related differences in ST-segment elevation myocardial infarction: A Portuguese multicenter national registry analysis, *Revista Portuguesa de Cardiologia*, <https://doi.org/10.1016/j.repc.2024.06.005>