

# ORIGINAL ARTICLE

# Revista Portuguesa de Cardiologia Portuguese Journal of Cardiology www.revportcardiol.org



# Development of a predictive nomogram for postoperative acute respiratory distress syndrome in Stanford type A aortic dissection patients: A retrospective study



Huanan Liu<sup>1</sup>, Hua Lu<sup>1</sup>, Zhaoming Lin, Xiaoshen Zhang\*

Department of Cardiovascular Surgery, The First Affiliated Hospital of Jinan University, Guangzhou, China

Received 18 May 2024; accepted 17 October 2024 Available online 25 January 2025

#### **KEYWORDS** Abstract Acute respiratory Introduction and objectives: This retrospective study aimed to develop a nomogram to predict distress syndrome; the risk of postoperative acute respiratory distress syndrome (ARDS) in patients with Stanford Type A aortic type A acute aortic dissection. dissection; Methods: The study included patients who underwent surgical repair for Stanford type A acute aortic dissection between January 2020 and December 2023. Demographic data, surgical details. Risk factor: Nomogram intraoperative information, and postoperative outcomes were collected. Univariate logistic regression was used for preliminary predictor screening, and a multivariate logistic regression model was constructed and presented as a nomogram. The nomogram's performance was evaluated using the area under the receiver operating characteristics (ROC) curve, calibration plots, and decision curve analysis (DCA). Internal validation was performed using bootstrap resampling. Results: The study included 142 patients, 41 (28.873%) of whom developed ARDS postoperatively. Multivariate logistic regression identified body mass index (BMI), postoperative procalcitonin (PCT), cardiopulmonary bypass (CPB) time, and low albumin as independent risk factors for postoperative ARDS in type A acute aortic dissection patients. These factors were used to develop the nomogram, which demonstrated good predictive performance with an area under the ROC curve of 0.809 (95% confidence interval: 0.721-0.881). The nomogram was successfully validated by calibration plots and DCA.

\* Corresponding author.

https://doi.org/10.1016/j.repc.2024.10.006

E-mail address: xsh.zhang@hotmail.com (X. Zhang).

<sup>&</sup>lt;sup>1</sup> These authors contributed equally to this work and share first authorship.

<sup>0870-2551/© 2025</sup> Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

*Conclusions:* BMI, PCT, CPB time, and low albumin are independent risk factors for postoperative ARDS in type A acute aortic dissection patients. The constructed nomogram provides an effective tool for predicting the risk of ARDS, aiding in the prevention and management of this complication in patients undergoing aortic surgery.

© 2025 Sociedade Portuguesa de Cardiologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

#### PALAVRAS-CHAVE

Síndrome do desconforto respiratório agudo; Disseção aórtica tipo A; Fator de risco; Nomograma

#### Nomograma para predição do risco de síndrome do desconforto respiratório agudo pós-operatório em pacientes com disseção aórtica tipo A: uma análise retrospectiva

#### Resumo

*Introdução e objetivos*: Este estudo retrospetivo teve como objetivo desenvolver um nomograma para prever o risco de síndrome do desconforto respiratório agudo (SDRA) no pós-operatório de pacientes com disseção aguda da aorta tipo A de Stanford (TAAD).

*Métodos:* Foram incluídos pacientes submetidos à correção cirúrgica para disseção aguda da aorta tipo A de Stanford entre janeiro de 2020 e dezembro de 2023. Foram coletados dados demográficos, detalhes cirúrgicos, informações intraoperatórias e desfechos pós-operatórios. A regressão logística univariada foi utilizada para triagem preliminar dos preditores, seguida pela construção de um modelo multivariado de regressão logística, apresentado como nomograma. O desempenho do nomograma foi avaliado por meio da área sob a curva ROC (Receiver Operating Characteristic), gráficos de calibração e análise da curva de decisão (DCA). A validação interna foi realizada por meio de reamostragem *bootstrap*.

*Resultados:* Foram incluídos 142 pacientes, dos quais 41 (28,9%) desenvolveram SDRA no pós-operatório. A regressão logística multivariada identificou o índice de massa corporal (IMC), a procalcitonina pós-operatória (PCT), o tempo de circulação extracorpórea (CEC) e a baixa albumina como fatores de risco independentes para SDRA pós-operatória em pacientes com TAAD. Esses fatores foram utilizados para desenvolver o nomograma, que demonstrou bom desempenho preditivo com uma área sob a curva ROC de 0,809 (intervalo de confiança de 95%: 0,721-0,881). O nomograma foi validado com sucesso por meio de curvas de calibração e DCA. *Conclusões:* IMC, PCT, tempo de CEC e baixa albumina são fatores de risco independentes para SDRA pós-operatória em pacientes com TAAD. O nomograma desenvolvido fornece uma ferramenta eficaz para prever o risco de SDRA, auxiliando na prevenção e no manejo dessa complicação em pacientes submetidos à cirurgia aórtica.

© 2025 Sociedade Portuguesa de Cardiologia. Publicado por Elsevier España, S.L.U. Este é um artigo Open Access sob uma licença CC BY-NC-ND (http://creativecommons.org/licenses/by-nc-nd/4.0/).

### Introduction

Acute respiratory distress syndrome (ARDS) is a clinical syndrome characterized by refractory hypoxemia, radiographic severity, and respiratory system compliance. This syndrome is usually associated with sepsis, shock, aspiration, and trauma. Cardiac surgery involving cardiopulmonary bypass (CPB) has been reported to result in ARDS in 0.4–20% of patients, with mortality rates reaching as high as 70–80%.<sup>1–3</sup> Beyond mortality, compromised long-term health status and increased medical expenditures could not be ignored contributing to this respiratory failure. Thus, every attempt to decrease the incidence of ARDS should be embraced.

Type A aortic dissection, which involves the ascending aorta irrespective of the site of the intimal tear according to the Stanford system, remains the most common aortic catastrophe despite recent advances in diagnosis and treatment. Prompt surgical repair is required because of an extremely high mortality 22.7% within the first six hours, 33.3% within 12 hours, 50% within 24 hours, and 68.2% within the first two days if left untreated.<sup>4</sup> Acute lung injury (ALI) and its more severe form, ARDS, are lethal complications during perioperative management; aortic surgery accounts for 16% of ARDS cases.<sup>2</sup> ALI and hypoxemia were observed in more than half of type A acute aortic dissection patients, with ARDS occurring in 10–20% of the population.<sup>5,6</sup> Additionally, emergent surgery predisposed type A acute aortic dissection patients to higher risk of ARDS.<sup>7</sup> Thus, a better understanding of the independent risk factors for ARDS in type A acute aortic dissection is essential to improve clinical strategies and outcomes.

However, drawing conclusions is challenging because studies on postoperative ARDS have primarily included patients undergoing heterogeneous cardiac surgeries, such as coronary artery bypass grafting (CABG) and multiple valve surgery. Only one recent study reported an occurrence of 15.9% in ARDS following aortic dissection surgery.<sup>6</sup> More research is needed to identify pre- and perioperative risk factors for ARDS. Therefore, we firstly established and validated a predictive nomogram to calculate the probability of ARDS for individuals with type A acute aortic dissection, which might enhance the clinical decision-making.

## Objectives

The study aimed to identify independent risk factors and establish a numerical model for postoperative acute respiratory distress syndrome in type A aortic dissection patients.

## Methods

## Study participants and design

This was a single-center, retrospective, observational study. Clinical data were collected from the hospital database between January 2020 and December 2023. Eligible participants were over 18 years old. The exclusion criteria for this study were as follows: (1) death before surgery; (2) no surgery performed; (3) incomplete records exceeding 20%. One hundred forty-two clinical records were enrolled into this study.

This study was approved by the Institutional Review Board of the First Affiliated Hospital of Jinan University (No.: KY-2023-001). Informed consent was obtained from all patients (or their guardians). All personal information was de-identified prior to analysis. The study was conducted in accordance with the Declaration of Helsinki.

## Clinical data and definitions

Acute respiratory distress syndrome was defined in accordance with the 2012 Berlin definition as follows<sup>8</sup>: (1) hypoxemia, acute onset within first week of exposure to a known clinical insult or new or worsening respiratory symptoms; (2) bilateral opacities on chest image that cannot be fully explained by the presence of pleural effusions, lobar or lung collapse, or nodules; and (3) clinical outcomes not fully explained by fluid overload or heart failure. Imaging findings were reviewed independently by two qualified physicians, and a third one was consulted in cases of disagreement over the diagnosis.

Data on demographic and clinical characteristics, including age, gender, body mass index (BMI), mean arterial pressure, vital signs, and cardiac function as well as comorbidities such as smoking, cerebrovascular disease, hypertension, diabetes, end-stage renal disease, and coronary heart disease were extracted. Surgery details, including the procedure, blood product transfusion, duration of the operation, CPB, mechanical ventilation, and aortic clamping, were recorded. Preoperative laboratory test results, including leukocyte count, hemoglobin, albumin, cardiac troponin I, N-terminal B-type natriuretic peptide, fibrinogen, D-dimer, prothrombin time, and serum creatinine were also collected. Postoperatively, all patients were transferred to the intensive care unit (ICU), where chest X-rays were conducted daily for the first three days. Serum procalcitonin (PCT) levels were measured on the first postoperative day. Postoperative complications, including stroke, deep sternal wound infection, tracheostomy, disseminated intravascular coagulation, liver failure, acute renal failure, and parenternal nultrition, were documented during follow-up until discharge from hospital.

## Statistical analysis

Data were collected and analyzed using R software package (Version 4.2.1) and DCPM (V2.33 Jingding Medical Technology Co., Ltd). Variables with a p-value  $\leq 0.05$  in the univariate logistic regression were considered potential predictors for postoperative ARDS and entered into a multivariate logical regression analysis with backward elimination, using the ''glm'' package in R. Results were presented as odds ratios (OR) with 95% confidence intervals (CIs). The prediction model established through multivariate logistic regression was visualized with a nomogram using the "rms" package in R. Model performance was evaluated using receiver operating characteristic (ROC) curve analysis and calibration plots, with a bootstrap resampling method (bootstrap=500). The area under the working characteristic curve (AUC) was used to assess model discrimination, while calibration was estimated using Hosmer-Lemeshow test. Clinical utility was assessed by decision curve analysis (DCA) combined with a cutoff value. All tests were two-tailed tests and p-values <0.05 were considered statistically significant. Variables with missing data exceeding 20% were excluded to ensure data reliability. A flow diagram depicting the study process is provided (Figure 1).

## Results

## Characteristics of the study population

A total of 172 patients diagnosed with Stanford type A aortic dissection preliminarily fulfilled the entry criteria for this study. Of these, 30 were excluded due to the following reasons: 15 cases died before surgery, 11 cases did not undergo surgery, and four cases had incomplete records. One hundred forty-two adult patients were enrolled, with a mean age of  $52.387 \pm 11.587$  years; 87.324%were male (18 females and 124 males). Among the enrolled patients, 41 (28.873%) developed postoperative ARDS. The study population exhibited various comorbidities: a history of smoking was observed in 40 patients (28.169%), cerebrovascular disease in 13 patients (9.155%), hypertension in 95 patients (66.901%), diabetes in six patients (4.225%), endstage renal disease in seven patients (4.930%) and coronary heart disease in four patients (2.817%). According to the New York Heart Association (NYHA) classification, nine patients (6.338%) were considered class II, 43 patients (30.282%) were considered class III, and 90 patients (63.380%) were considered class IV.

All patients underwent aortic repair surgery through median sternotomy, with CPB established through femoral artery and right axillary artery cannulation. Thirty-four patients (23.944%) received Bentall surgery, which involved replacing the aortic valve with a mechanical valve; nine



Figure 1 Study design. ARDS: acute respiratory distress syndrome.

patients (6.338%) underwent CABG, and two cases (1.401%) had mitral and/or tricuspid valve repair. The mean durations of the operations, CPB, aortic clamping, and mechanical ventilation were 473.754 $\pm$ 99.719 minutes, 237.831 $\pm$ 65.652 minutes, 129.690 $\pm$ 34.941 minutes, and 45.500 (20.500, 107.750) hours, respectively. The average length of ICU stay was 9.620 $\pm$ 9.238 days. The all-cause mortality rate in hospital was 7.745% (11 cases).

#### Risk factors and the prediction model

Patients in the ARDS group had significantly higher BMI ( $26.935\pm4.650$  vs.  $25.034\pm3.782$  kg/m<sup>2</sup>, p=0.012), leukocyte counts ( $13.165\pm4.448$  vs.  $11.417\pm4.228\times10^9$ , p=0.030), PCT [21.960 (4.557, 43.000) vs. 4.643 (2.408, 12.855) ng/mL, p<0.001], as well as longer durations of operation ( $505.122\pm105.575$  vs.  $461.020\pm94.843$  min, p=0.016) and CPB time ( $268.244\pm69.305$  vs.  $225.485\pm60.191$ 

min, p<0.001) compared to patients in the non-ARDS group. A higher proportion of patients who underwent CABG [6 (14.634%) vs. 3 (2.970%), p=0.027] was observed in the ARDS group. Additionally, albumin levels were notably decreased in ARDS patients (34.341 $\pm$ 5.376 vs. 36.629 $\pm$ 4.456 g/L, p=0.010).

Univariate analysis identified several potential risk factors for postoperative ARDS (Table 1): BMI (OR: 1.120; 95% CI: 1.022–1.228), leukocyte (1.096; 1.007–1.192), albumin (0.907; 0.840–0.980), PCT (1.017; 1.005–1.029), CABG (5.600; 1.329–23.604), operation time (1.005; 1.001–1.008), and CPB time (1.010; 1.004–1.016). Variables with a p-value  $\leq$ 0.05 were entered into a multivariate logical regression analysis. The results of the multivariate analysis, illustrated in Table 2 and visualized in a forest plot (Figure 2), identified four predictors of postoperative ARDS: BMI (OR: 1.261; 95% CI: 1.121–1.444), albumin (0.850; 0.762–0.937), PCT (1.023; 1.011–1.038), and CPB time (1.008; 1.002–1.015).

## Table 1 Univariate analysis of variables related to postoperative ARDS in TAAD patients.

Variables	OR	95% CI	р	Variables	OR	95% CI	р
Demographics				Operations			
Age	1.020	[0.988, 1.053]	0.231	Bentall			
Gender				No	Ref.		
F	Ref.			Yes	0.447	[0.170, 1.178]	0.104
Μ	2.209	[0.604, 8.083]	0.231	CABG		- / -	
ВМІ	1.120	[1.022, 1.228]	0.016*	No	Ref.		
Vital signs				Yes	5.600	[1.329, 23.604]	0.019*
Heart rate	1.006	[0.985, 1.028]	0.568	Operation time	1.005	[1.001, 1.008]	0.019*
MAP	0.977	[0.954, 1.001]	0.057	CPB time	1.010	[1.004, 1.016]	0.001*
Cardiac anatomy and function				Aortic cross	1.010	[0.999, 1.021]	0.058
				clamping time			
NYHA class				Mechanical	1.003	[0.999, 1.006]	0.141
				ventilation time		- / -	
II	Ref.			Blood transfusion			
III	1.556	[0.167, 14.480]	0.698	PRBC	1.084	[0.955, 1.229]	0.212
IV	4.632	[0.555, 38.685]	0.157	FFP	1.001	[0.999, 1.003]	0.062
LVEF	0.989	[0.949, 1.031]	0.601	Cryoprecipitate	0.823	[0.649, 1.044]	0.108
LA diameter	1.004	[0.943, 1.069]	0.898	PLT	0.941	[0.562, 1.575]	0.817
LV diameter	0.958	[0.904, 1.015]	0.148	Pre. $PaO_2/FiO_2$	0.998	[0.994, 1.002]	0.280
RA diameter	0.949	[0.888, 1.013]	0.115	Post. $PaO_2/FiO_2$	0.996	[0.991, 1.001]	0.117
		[]		in day 1		[]	
RV diameter	0.976	[0.879, 1.084]	0.646	Post. PaO <sub>2</sub> /FiO <sub>2</sub>	0.999	[0.995, 1.003]	0.452
		[]		in day 3		[]	
Pulmonary artery systolic pressure	1.045	[0.980, 1.113]	0.180	Drainage	1.000	[0,999, 1,002]	0.683
, a		[01/00, 1110]		volume in first 24h		[01777, 1000-]	
Diameter of the aortic root	0.993	[0.937, 1.051]	0.796	Postoperative	0.997	[0.966, 1.018]	0.539
brameter of the dortheroot	01775	[0.757, 1.051]	0.770	hospital stav	0.772	[01700, 11010]	0.557
Laboratory tests				ICU stav	1.001	[0.963, 1.042]	0.947
Leukocyte	1.096	[1 007 1 192]	0.033*	Postoperative		[01700, 110.2]	
Leakocyte		[]	0.000	complications			
Hemoglobin	0.997	[0.975, 1.020]	0.795	Cerebral			
hemogeoni	0.777	[01770, 11020]	0.775	vascular accident			
Albumin	0 907	[0 840 0 980]	0.013*	No	Ref		
NT-pro BNP	0.999	[0 999 1 000]	0.013	Yes	0.815	[0 370 1 795]	0 612
Serum creatinine	1 001	[0 998 1 004]	0.668	Deen sternal	0.015	[0.570, 1.775]	0.012
Ser am er caemme		[0.770, 1.001]	0.000	wound infection			
cTNI	1 020	[0 993 1 047]	0 146	No	Ref		
D-dimer	1 000	[1 000 1 000]	0 335	Yes	0 985	[0 183 5 291]	0 986
Fibringen	0.833	[0 649 1 067]	0.333	Tracheostomy	0.705	[0.105, 5.271]	0.700
PT	1 040	[0 945 1 143]	0 474	No	Ref		
PCT	1.017	[1 005 1 029]	0.004*	Yes	0 480	[0 054 4 240]	0 509
Comorbidities		[11005, 11027]	0.001	DIC	01100		0.507
Smoking				No	Ref		
No	Ref			Yes	2 667	[0 729 9 760]	0 138
Ves	1 079	[0 484 2 405]	0 853	l iver failure	2.007	[0.727, 7.700]	0.150
Cerebrovascular disease	1.077	[0.404, 2.405]	0.055	No	Rof		
No	Rof			Yes	1 460	[0 691 3 083]	0 322
Vos	0 718	[0 187 2 756]	0.630	Acuto ronal	1.400	[0.071, 5.005]	0.322
103	0.710	[0.107, 2.750]	0.050	failure			
Hupertension				No	Rof		
No	Pof			Vor	1 455	[0 707 2 440]	0 177
Vec	1 282	[0 583 2 817]	0 537	Darenteral	1.055	[0.777, 3.440]	0.177
103	1.202	[0.303, 2.017]	0.337	nutrition			

#### Table 1 (Continued)

Variables	OR	95% CI	р	Variables	OR	95% CI	р
Diabetes				No	Ref.		
No	Ref.			Yes	1.380	[0.649, 2.935]	0.403
Yes	2.579	[0.499, 13.342]	0.259				
End-stage renal disease							
No	Ref.						
Yes	0.985	[0.183, 5.291]	0.986				
Coronary heart disease							
No	Ref.						
Yes	0.817	[0.083, 8.088]	0.863				

ARDS: acute respiratory distress syndrome; BMI: body mass index; CABG: coronary artery bypass grafting; CI: confidential interval; CPB: cardiopulmonary bypass; cTNI: cardiac troponin I; DIC: disseminated intravascular coagulation; FFP: fresh frozen plasma; FiO<sub>2</sub>: fraction of inspired oxygen; ICU: intensive care unit; LA: left atrium; LV: left ventricle; LVEF: left ventricular ejection fraction; MAP: mean arterial pressure; NT-pro BNP: N-terminal B-type natriuretic peptide; NYHA: New York Heart Association function classification; OR: odds ratio; PaO<sub>2</sub>: partial pressure of oxygen in the arterial blood; PCT: procalcitonin; PLT: platelet; PRBC: packed red blood cell; PT: prothrombin time; RA: right atrium; RV: right ventricle; TAAD: type A aortic dissection.

<sup>\*</sup> Significant at a p value  $\leq 0.05$ .

Table 2 Multivariate logistic regression analysis of independent risk factors for postoperative ARDS in TAAD patients.

Variables	OR	95% CI	p-Value
BMI	1.261	[1.121, 1.444]	<0.001
Albumin	0.850	[0.762, 0.937]	0.002
PCT	1.023	[1.011, 1.038]	<0.001
CPB time	1.008	[1.002, 1.015]	0.014

BMI: body mass index; CPB: cardiopulmonary bypass; PCT: procalcitonin.



**Figure 2** Forest plot according to multivariate regression analysis. BMI: body mass index; CI: confidential interval; CPB: cardiopulmonary bypass; OR: odds ratio; PCT: procalcitonin.

A nomogram was developed incorporating BMI, albumin, PCT, and CPB time to predict the risk of postoperative ARDS in type A acute aortic dissection patients (Figure 3). Each predictor was assigned a score range based on its value, and the total score, derived from summing the points for each factor, can be used to estimate the probability of ARDS.

#### Model validation and net benefit prediction

The predictive model for estimating the occurrence of ARDS after aortic surgery demonstrated stable performance, as validated by bootstrapping with 500 replications (Figure 4). The pooled area under the ROC curve was 0.809 (95% CI:



Figure 3 Nomogram for the risk of postoperative ARDS in TAAD patients. TAAD: type A aortic dissection.



**Figure 4** Bootstrapped ROC curve of the nomogram for postoperative ARDS in TAAD patients (AUC: 0.809, 95% CI: 0.721–0.881, bootstrap replicates=500).

0.721–0.881). The optimal cutoff value for predicting postoperative ARDS was 0.343, providing a high specificity of 0.830 and a sensitivity of 0.659. The Hosmer–Lemeshow test ( $\chi^2$ =3.664, p=0.932) indicated a good concordance between the predicted probabilities and the observed postoperative ARDS in type A acute aortic dissection patients (Figure 5). Boosted DCA curve displayed clinical benefit in Figure 6, with a threshold probability range of 0.01–0.95. The horizontal line represents the scenario in which no patients develop ARDS and thus no benefit from intervention. The diagonal line represents the scenario in which all patients develop ARDS, showing potential benefit from treatment. Therefore, the DCA confirms the clinical practicality of the prediction model within this range.

## Discussion

This study is the first to develop a nomogram of postoperative ARDS in type A acute aortic dissection patients, which illustrates the regression model in an accessible way and thus simplifies the risk assessment. We analyzed clinical characteristics, laboratory values and postoperative outcomes to identify the most relevant variables for multivariate regression analysis. Four independent predictors for postoperative ARDS were identified: BMI, albumin, PCT, and CPB time.

Postoperative ARDS carries a high mortality rate, ranging from 10% to 70%, resulting from the differences in ARDS definitions, types of surgeries, and study populations.<sup>1,6,9-11</sup> Previous definitions of ARDS, such as the Murray's Lung Injury Score from 1988, the American-European Consensus Conference definition (AECC) from 1994, and the Berlin definition from 2012, offer varying criteria for diagnosis and intervention.<sup>8,12,13</sup> The Berlin definition, which is the most recent one, provided practical guidelines for early recognition and intervention of ARDS. However, the application of the Berlin definition to type A acute aortic dissection patients specifically has not been thoroughly investigated. Studies on ARDS following cardiac surgery often included a mix of procedures involving CPB, such as valvular surgery and CABG. These varied combinations can lead to different incidence and mortality rates. For instance, a study reported that 8.1% of 457 patients undergoing isolated valvular heart surgery developed postoperative ARDS, with a mortality rate of 29.7%.<sup>10</sup> In contrast, a broader study including CABG, valve, and combined procedures found mortality of 68.4%.<sup>11</sup> Our study observed an ARDS incidence of 28.873% in type A acute aortic dissection patients, which was slightly higher than reported in some studies, possibly due to the high proportion of patients classified as NYHA class IV.<sup>6</sup>



Figure 5 Calibration plot of the nomogram for postoperative ARDS in TAAD patients (bootstrap replicates=500).



Figure 6 DCA curve of the nomogram for postoperative ARDS in TAAD patients (bootstrap replicates=500).

The pathophysiology of ARDS following CPB has not been totally understood. The prevailing hypothesis involves activation of the inflammatory cascade, including polymorphonuclear leukocyte and cytokine activation, which increases pulmonary permeability and vascular resistance, leading to impaired gas exchange.<sup>14</sup> Our findings indicate that CPB time is a significant risk factor in ARDS. Besides, protamine-heparine complexes may trigger immune reactions by releasing anaphylatoxins such as C4a and C3a, potentially increasing ARDS risk.<sup>1</sup> Patients with ARDS were reported to have lower preoperative albumin, which was similar in our study, and higher plasminogen activator inhibitor-1, interleukin-8 and surfactant protein-D.<sup>15</sup> Upregulation of tumor necrosis factor-alpha and elevation of serum endocan are also believed to get involved in the early phase of ARDS.<sup>16,17</sup> Albumin in the extracorporeal prime can preserve platelet count and reduce chest tube drainage by passivating the synthetic surfaces of the extracorporeal circuit.<sup>18</sup> Elevated PCT levels were also associated with ARDS in our study, suggesting it could serve as a potential diagnostic biomarker.<sup>19</sup> Timely infection control and interruption of inflammatory processes are crucial for ARDS prevention. Corticosteroids have been shown to improve cardiopulmonary function within three to seven days and alter the ARDS course, with early treatment improving ICU and ventilator outcomes with minimal safety concerns.<sup>20</sup>

Transfusion-related complications, including transfusionrelated acute lung injury and transfusion-associated circulatory overload, are significant concerns. Multiple transfusions have been linked to ARDS risk, with packed red blood cell transfusions exhibiting a dose-dependent effect on pulmonary capillary permeability.<sup>21</sup> In a large observational cohort of CABG patients (11963), red blood cell transfusion was associated with higher mortality, renal failure, prolonged ventilatory support.<sup>22</sup> A retrospective nested case-control study demonstrated extended durations of CPB, deep hypothermic circulatory arrest, ICU stay, and larger perioperative transfusions volumes (red blood cell, platelet and fresh frozen plasma) in ARDS group of type A acute aortic dissection population.<sup>23</sup> Although our study did not identify blood product transfusion as an independent predictor of ARDS, possibly due to sample size limitations and the urgent nature of aortic surgery, meticulous hemostasis and restrictive transfusion strategies could mitigate ARDS risk. A restrictive transfusion strategy targeting a hematocrit of 24% has been shown to be as safe as a liberal strategy in elective cardiac surgery, though caution is needed for elderly patients due to cardiovascular risks.<sup>24,25</sup>

Obesity's role in ARDS remains debated and is known as the ''obesity paradox''. While excess body weight has not consistently been linked to increased mortality in acute lung injury cases,<sup>26,27</sup> it is associated with longer hospital stays.<sup>28,29</sup> A study of 5023 patients undergoing heart surgery with CPB found that obese patients were at higher risk for postoperative hypoxia, but were somewhat protected against adverse effects from hemodilution and transfusions.<sup>30</sup> Our study confirmed that BMI is an independent risk factor of ARDS after aortic surgery. A case-control study identified a BMI cutoff of 24.78 kg/m<sup>2</sup>, with ARDS likelihood increasing above this threshold.<sup>31</sup> While BMI is challenging to modify in emergency settings, weight control strategies could benefit postoperative outcomes.

This study has limitations, including the inability to split the data into training and validation sets due to a limited sample size. Consequently, model validation was performed using bootstrap methods. Additionally, among data from a monocenter there may be some degree of bias. Further research with multicenter data and larger sample sizes is needed to enhance the validity and generalizability of these findings.

## Conclusion

We established a straightforward predictive nomogram to estimate the probability of ARDS in patients with type A acute aortic dissection by incorporating four perioperative variables, including BMI, albumin, procalcitonin and CPB time. The accuracy and stability of the model were validated internally using the AUC, calibration plot, and DCA. This nomogram has the potential to serve as a valuable prognostic tool for assessing the risk of ARDS in type A acute aortic dissection patients.

### Funding

None declared.

### **Conflicts of interest**

None declared.

#### References

- 1. Milot J, Perron J, Lacasse Y, et al. Incidence and predictors of ARDS after cardiac surgery. Chest. 2001;119:884–8.
- Gajic O, Dabbagh O, Park PK, et al. Early identification of patients at risk of acute lung injury: evaluation of lung injury prediction score in a multicenter cohort study. Am J Respir Crit Care Med. 2011;183:462–70.
- Rong LQ, Di Franco A, Gaudino M. Acute respiratory distress syndrome after cardiac surgery. J Thorac Dis. 2016;8:E1177–86.
- Mészáros I, Mórocz J, Szlávi J, et al. Epidemiology and clinicopathology of aortic dissection. Chest. 2000;117:1271–8.
- Pan X, Lu J, Cheng W, et al. Independent factors related to preoperative acute lung injury in 130 adults undergoing Stanford type-A acute aortic dissection surgery: a single-center crosssectional clinical study. J Thorac Dis. 2018;10:4413–23.
- Su IL, Wu VC, Chou AH, et al. Risk factor analysis of postoperative acute respiratory distress syndrome after type A aortic dissection repair surgery. Medicine (Baltimore). 2019;98:e16303.
- 7. Fernández-Pérez ER, Sprung J, Afessa B, et al. Intraoperative ventilator settings and acute lung injury after elective surgery: a nested case control study. Thorax. 2009;64:121–7.
- Ranieri VM, Rubenfeld GD, Thompson BT, et al. Acute respiratory distress syndrome: the Berlin definition. JAMA. 2012;307:2526-33.
- 9. Kogan A, Preisman S, Levin S, et al. Adult respiratory distress syndrome following cardiac surgery. J Card Surg. 2014;29:41–6.
- **10.** Chen SW, Chang CH, Chu PH, et al. Risk factor analysis of postoperative acute respiratory distress syndrome in valvular heart surgery. J Crit Care. 2016;31:139–43.
- Christenson JT, Aeberhard JM, Badel P, et al. Adult respiratory distress syndrome after cardiac surgery. Cardiovasc Surg. 1996;4:15–21.
- Murray JF, Matthay MA, Luce JM, et al. An expanded definition of the adult respiratory distress syndrome. Am Rev Respir Dis. 1988;138:720-3.
- Bernard GR, Artigas A, Brigham KL, et al. The American-European Consensus Conference on ARDS, definitions, mechanisms, relevant outcomes, and clinical trial coordination. Am J Respir Crit Care Med. 1994;149 Pt 1:818–24.
- 14. Yousefshahi F, Samadi E, Paknejad O, et al. Prevalence and risk factors of hypoxemia after coronary artery bypass grafting: the time to change our conceptions. J Tehran Heart Cent. 2019;14:74–80.
- **15.** Yadav H, Bartley A, Keating S, et al. Evolution of validated biomarkers and intraoperative parameters in the development of postoperative ARDS. Respir Care. 2018;63:1331–40.

- 16. Li T, Luo N, Du L, et al. Early and marked up-regulation of TNFα in acute respiratory distress syndrome after cardiopulmonary bypass. Front Med. 2012;6:296–301.
- **17.** Bouglé A, Allain PA, Favard S, et al. Postoperative serum levels of Endocan are associated with the duration of norepinephrine support after coronary artery bypass surgery. Anaesth Crit Care Pain Med. 2018;37:565–70.
- Beney KC. Human albumin in extracorporeal prime: effect on platelet function and bleeding. Perfusion. 2013;28:536–40.
- Cheng ZB, Chen H. Higher incidence of acute respiratory distress syndrome in cardiac surgical patients with elevated serum procalcitonin concentration: a prospective cohort study. Eur J Med Res. 2020;25:11.
- Steinberg KP, Hudson LD, Goodman RB, et al. Efficacy and safety of corticosteroids for persistent acute respiratory distress syndrome. N Engl J Med. 2006;354:1671–84.
- 21. Vlaar AP, Cornet AD, Hofstra JJ, et al. The effect of blood transfusion on pulmonary permeability in cardiac surgery patients: a prospective multicenter cohort study. Transfusion. 2012;52:82–90.
- 22. Koch CG, Li L, Duncan AI, et al. Morbidity and mortality risk associated with red blood cell and blood-component transfusion in isolated coronary artery bypass grafting. Crit Care Med. 2006;34:1608–16.
- 23. Chen MF, Chen LW, Cao H, et al. Analysis of risk factors for and the prognosis of postoperative acute respiratory distress syndrome in patients with Stanford type A aortic dissection. J Thorac Dis. 2016;8:2862–71.

- 24. Hajjar LA, Vincent JL, Galas FR, et al. Transfusion requirements after cardiac surgery: the TRACS randomized controlled trial. JAMA. 2010;304:1559–67.
- **25.** Nakamura RE, Vincent JL, Fukushima JT, et al. A liberal strategy of red blood cell transfusion reduces cardiogenic shock in elderly patients undergoing cardiac surgery. J Thorac Cardiovasc Surg. 2015;150:1314–20.
- **26.** Newell MA, Bard MR, Goettler CE, et al. Body mass index and outcomes in critically injured blunt trauma patients: weighing the impact. J Am Coll Surg. 2007;204:1056–61.
- 27. O'Brien JJ, Welsh CH, Fish RH, et al. Excess body weight is not independently associated with outcome in mechanically ventilated patients with acute lung injury. Ann Intern Med. 2004;140:338–45.
- **28.** Gong MN, Bajwa EK, Thompson BT, et al. Body mass index is associated with the development of acute respiratory distress syndrome. Thorax. 2010;65:44–50.
- **29.** Dossett LA, Heffernan D, Lightfoot M, et al. Obesity and pulmonary complications in critically injured adults. Chest. 2008;134:974–80.
- Ranucci M, Ballotta A, La Rovere MT, et al. Postoperative hypoxia and length of intensive care unit stay after cardiac surgery: the underweight paradox? PLoS One. 2014;9:e93992.
- Liu Y, Song M, Huang L, et al. "Obesity paradox" in acute respiratory distress syndrome among patients undergoing cardiac surgery: a retrospective study. Med Sci Monit. 2021;27:e931808.