

IMAGE IN CARDIOLOGY

Pulmonary artery sling: An incidental finding**Sling da artéria pulmonar: um achado acidental**

Joana O. Miranda*, José Monterroso, Jorge Moreira, Maria João Baptista



Serviço de Cardiologia Pediátrica, Centro Hospitalar São João, Porto, Portugal

Received 2 January 2014; accepted 27 January 2014

Available online 20 August 2014

A previously healthy one-month infant was referred to our institution due to a heart murmur. The transthoracic echocardiogram showed the pulmonary trunk giving rise to the right pulmonary artery (RPA), but the left pulmonary artery (LPA) was not identified in its usual location. Instead the LPA was arising from the RPA and coursing leftward behind the trachea, with a turbulent flow (peak velocity 1.5 m/s) (Video 1). An associated ostium secundum atrial septal defect was diagnosed. The suspected diagnosis of pulmonary artery sling was confirmed by computed tomographic angiography. The LPA was seen originating from the posterior aspect of the RPA and coursing over the right mainstem bronchus and then from right to left, posterior to the trachea and anterior to the esophagus, to reach the hilum of the left lung, and causing a mild compression of the origin of the LPA (Figure 1). Besides the pulmonary artery sling and the atrial septal defect, a partial anomalous pulmonary venous return was identified, with the left superior pulmonary vein draining in the left brachiocephalic vein (Figure 2).

The anomalous origin of the LPA known as pulmonary artery sling is a rare form of vascular ring. In the vast majority of patients it is associated with respiratory symptoms in the first year of life, due to tracheal stenosis, and with a




Figure 1 Computed tomographic angiography reveals anomalous origin of the LPA from the posterior aspect of the RPA. The aberrant LPA (black arrow) runs behind the trachea as it courses to the left pulmonary hilum, causing a mild compression of the origin of the LPA.

high mortality rate if there is no surgical intervention. In our case, the heart murmur led to early medical attention. The findings on echocardiography and computed tomographic angiography allowed an early non-invasive diagnosis, avoiding a potentially severe and life-threatening acute presentation.

* Corresponding author.

E-mail address: joanam@gmail.com (J.O. Miranda).



Figure 2 Computed tomographic angiography reveals partial anomalous pulmonary venous return consisting of left superior pulmonary vein (white arrow) draining in the left brachiocephalic vein.

Conflict of interest

The authors have no conflict of interest to declare.

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that they have followed the protocols of their work center on the publication of patient data.

Right to privacy and informed consent. The authors have obtained the written informed consent of the patients or subjects mentioned in the article. The corresponding author is in possession of this document.

Appendix A. Supplementary data

Supplementary data associated with this article can be found in the online version, at [doi:10.1016/j.repc.2014.01.022](https://doi.org/10.1016/j.repc.2014.01.022).