



ORIGINAL ARTICLE

The genetic variant C825T of the beta 3 subunit of G protein is associated with hypertension in a Portuguese population[☆]



Ana Célia Sousa^{a,*}, Roberto Palma dos Reis^b, Andreia Pereira^a, Sofia Borges^a, Sara Gouveia^a, Adelaide Spínola^a, Ana Isabel Freitas^{a,c}, Graça Guerra^{a,c}, Teresa Góis^a, Mariana Rodrigues^a, Eva Henriques^a, Ilídio Ornelas^a, Carolina Freitas^a, Décio Pereira^a, António Brehm^c, Maria Isabel Mendonça^a

^a *Unidade de Investigação, Hospital Doutor Nélio Mendonça, Funchal, Portugal*

^b *Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisboa, Portugal*

^c *Laboratório de Genética Humana, Universidade da Madeira, Funchal, Portugal*

Received 13 March 2017; accepted 24 September 2017

Available online 1 June 2018

KEYWORDS

Case-control study;
Hypertension;
G protein;
Polymorphisms;
GNB3

Abstract

Introduction: Hypertension is an important public health problem, affecting about 25% of the adult population worldwide.¹ Genetic and environmental factors contribute to its pathogenesis. The T allele of the C825T polymorphism of the beta 3 subunit of G protein (rs5443) leads to the production of a truncated variant that enhances intracellular signaling and may interfere with the regulation of blood pressure. This genetic variant has been described as a risk factor for hypertension, although study results are controversial.

Objective: The objective of this study was to analyze the association of the C825T polymorphism of the *GNB3* gene with the occurrence of hypertension in a Portuguese population from the Madeira archipelago.

Methods: A case-control study was performed with 1641 Caucasian individuals (mean age 50.6±8.1 years), 848 with hypertension and 793 controls. Blood was collected from all participants for biochemical and genetic analysis, including genotyping of the C825T polymorphism. Logistic regression analysis was performed to determine which variables were significantly associated with the onset of hypertension. Statistical analyses were performed using IBM SPSS version 19.0 and p-values <0.05 were considered statistically significant.

[☆] Please cite this article as: Sousa AC, Palma dos Reis R, Pereira A, Borges S, Gouveia S, Spínola A, et al. A variante genética c825t da subunidade β3 da proteína G associa-se com a hipertensão arterial numa população portuguesa. Rev Port Cardiol. 2018;37:499–507.

* Corresponding author.

E-mail address: dep.card@sesaram.pt (A.C. Sousa).

PALAVRAS-CHAVE

Estudo caso-controlo;
Hipertensão arterial;
Polimorfismos;
Proteína G;
GN β 3

Results: In our study, there was a significant association between the C825T polymorphism of the *GNB3* gene and the occurrence of hypertension (odds ratio 1.275; 95% confidence interval 1.042-1.559; $p=0.018$) in the dominant model, after multivariate analysis.

Conclusion: We conclude that the C825T polymorphism of the beta 3 subunit of G protein is significantly and independently associated with the occurrence of hypertension in the study population.

© 2018 Published by Elsevier España, S.L.U. on behalf of Sociedade Portuguesa de Cardiologia.

A variante genética c825t da subunidade β 3 da proteína G associa-se com a hipertensão arterial numa população portuguesa

Resumo

Introdução: A hipertensão arterial é um problema de Saúde Pública, afeta 25% da população adulta mundial. Fatores genéticos e ambientais contribuem para a sua patogénese. O polimorfismo C825T da subunidade β 3 da Proteína G (rs5443) favorece a produção de uma variante alternativa, truncada, que facilita a sinalização intracelular, pode interferir na regulação da pressão arterial. Essa variante genética tem sido descrita como um fator de risco para a hipertensão arterial, com resultados controversos.

Objetivo: Avaliar a associação do polimorfismo C825T do gene GN β 3 com o aparecimento de hipertensão arterial, numa população portuguesa do Arquipélago da Madeira.

Métodos: Com uma amostra de 1.641 indivíduos (média de $50,6 \pm 8,1$ anos), fizemos um estudo caso-controlo com 848 indivíduos com hipertensão arterial e 793 controlos, ajustados para o sexo e a idade. Todos os participantes colheram sangue para análises bioquímicas e foram genotipados para o polimorfismo C825T. Foi feita uma regressão logística para ver quais as variáveis que se relacionam com a hipertensão arterial. A análise dos dados foi feita com o *software* estatístico SPSS, versão 19.0. Usamos como limiar de significância o valor de $p < 0,05$.

Resultados: Encontramos uma associação significativa entre o polimorfismo C825T e o aparecimento de hipertensão arterial (*odds ratio* = 1,275; IC 95% (1,042–1,559); $p = 0,018$) no modelo dominante, após análise multivariada.

Conclusão: O polimorfismo C825T da subunidade β 3 da Proteína G está associado, de forma significativa e independente, com o aparecimento hipertensão arterial na nossa população.

© 2018 Publicado por Elsevier España, S.L.U. em nome de Sociedade Portuguesa de Cardiologia.

Introduction

Hypertension is a major public health problem, affecting around 25% of the adult population worldwide.¹ It is a risk factor for cardiovascular disease and progression to renal failure, independently of other cardiovascular risk factors.^{2,3} In the Portuguese adult population, its overall prevalence is around 42% (44.4% in men and 40.2% in women).⁴

In recent decades, many studies have been conducted to determine how genetics contributes to the development of hypertension. Around 20-40% of blood pressure variation is genetically determined,^{5,6} but the molecular genetics of HT is still unclear.

G proteins and their receptors play a crucial role in cell survival. These proteins are part of a superfamily of proteins that, when inactive, are bound to cell membrane receptors, and when activated migrate through the cytosol regulating multiple effectors including enzymes, ion channels, hormones, neurotransmitters and autocrine and paracrine

factors.⁷ They are thus signal transducers that enable intracellular events to be triggered by external stimuli.

Transduction of external stimuli via intracellular signals is clearly of considerable importance for perpetuating viable characteristics on an evolutionary scale. Given their wide-ranging roles, deficiency or altered expression of these important proteins can lead to more or less widespread metabolic disturbances.⁸

Heterotrimeric G proteins consist of three subunits, alpha, beta and gamma, coded by separate genes. The alpha subunit of each G protein is the most important: it is the subunit that interacts with the receptor, binding to guanosine triphosphate (GTP) and regulating effector systems.⁹ The beta and gamma subunits are non-covalently bound into the beta-gamma complex, which when bound to the alpha subunit forms the inactive state of the protein.¹⁰ In addition to anchoring the alpha subunit, the beta-gamma dimer influences certain cell processes^{11,12} and can modulate the activity of certain effectors.¹³ Due to their crucial role in the functioning of many cell types, genetic abnormalities in

G protein subunits may be involved in the etiology of a wide range of clinical conditions.

The gene that codes for the G protein beta 3 subunit is located on chromosome 12p13. The C825T single-nucleotide polymorphism of the beta 3 subunit of G protein (rs5443) is located in exon 10 of the *GNB3* gene, where cytosine (C) is replaced by thymidine (T) at nucleotide 825. The T allele of the C825T polymorphism results in a splicing variation in exon 9 in which nucleotides 498-620 are deleted,^{14,15} leading to the absence of amino acids 167 to 197 in the truncated protein (known as GNB3-s). This is a functional protein that enhances protein G-associated intracellular signaling.^{14,16} Since G-protein activation is the key event in intracellular signal transduction, the C825T polymorphism plays an important role in the etiology of a variety of disease processes,¹⁷ including hypertension,^{14,15,18-20} obesity,²¹ depression,²² and cardiovascular disease.²³ Furthermore, the 825T allele functions as a pharmacogenetic marker to predict responses to various drugs, such as diuretics, antidepressants, sildenafil and clonidine, as well as to angiotensin II and endothelin 1.¹⁷

However, the results of some studies have failed to demonstrate any association between this genetic variant and hypertension,²⁴⁻²⁶ while others support it. In view of this controversy, we decided to assess this association in a Portuguese population.

Objective

The objective of this study was to analyze the association of the C825T polymorphism of the *GNB3* gene (rs5443) with the occurrence of hypertension in a Portuguese population from the Madeira archipelago.

Methods

Study protocol and ethics approval

The study was approved by the ethics committee of the Health Service of the Autonomous Region of Madeira (SESARAM) and was conducted in accordance with the international ethics guidelines of the 2013 Declaration of Helsinki.

All participants gave their written informed consent, after an internist with more than 10 years of experience explained how blood would be collected for analysis and how DNA and relevant clinical data would be stored and reused. Blood samples for genetic analysis are kept in our hospital's research biobank in accordance with Articles 6, 15 and 16 of the UNESCO Universal Declaration on Bioethics and Human Rights.

Study population

This study was conducted on the Portuguese archipelago of Madeira, which has a population of around 300 000. Participants were Caucasian, born in Madeira, and their parents and grandparents were also Caucasians born in the archipelago. The study sample was thus a genetically homogeneous southern European population.

The sample consisted of 1641 individuals with a mean age of 50.6 ± 8.1 years, 848 (49.9%) male, who were selected sequentially from general and family medicine consultations in the Internal Medicine department of the Central Hospital of Funchal.

The population was divided into two groups, those with and those without hypertension. Differences between the groups in terms of gender and age were kept to a minimum.

Hypertension

At enrollment, subjects were defined as having hypertension if it had already been diagnosed, if they had been taking antihypertensive medication for more than three months, or if they were newly diagnosed with systolic (SBP) and/or diastolic blood pressure (DBP) $\geq 140/90$ mmHg on at least three occasions.²⁷

Individuals with secondary hypertension, such as those with primary hyperaldosteronism or clinically significant renal disease, were excluded, as were those taking medication for other indications that could affect blood pressure.

The normotensive control subjects had never taken antihypertensive medication and had SBP/DBP $< 140/90$ mmHg.

Blood pressure was measured in the right arm with the subject in a seated position after 10 minutes' rest, using a standard Welch Allyn sphygmomanometer. The mean of three readings taken at 2-min intervals was recorded.²⁷

Data collection

All participants answered a standardized questionnaire on age, gender, lifestyle, smoking, alcohol consumption, and personal and family history of cardiovascular events, particularly stroke, coronary artery disease and family history of hypertension.

Anthropometric parameters were assessed in all participants. Height was measured in cm and weight in kg, and body mass index (BMI) was calculated as kg/m^2 .²⁸ BMI was classified as normal ($< 25 \text{ kg}/\text{m}^2$), overweight ($25-29.9 \text{ kg}/\text{m}^2$) or obese ($\geq 30 \text{ kg}/\text{m}^2$).²⁸ Waist circumference was measured and considered high when > 94 cm in men and > 80 cm in women.²⁹ Waist-hip ratio was defined as high when > 0.9 in men and > 0.8 in women.²⁹

Pulse wave velocity (PWV) was determined by tonometry using a Complior device, as previously described.³⁰ PWV > 10 m/s was considered high. Diabetes was defined as fasting blood glucose ≥ 126 mg/dl and/or blood glucose ≥ 200 mg/dl and/or a history of treatment for diabetes.³¹

A sedentary lifestyle was defined as performing less than 150 min/week of moderate to vigorous physical activity.²⁹

Individuals who smoked at enrollment in the study were considered smokers and were classified according to the number of cigarettes smoked per day: 1-10, 10-20 or more than 20. Individuals who consumed alcohol at enrollment were considered to be drinkers. Alcohol consumption was quantified in g per week and was considered excessive when > 70 g/week in men and > 50 g/week in women.

Laboratory tests

Blood samples were collected after 14-16 hours of fasting and plasma was prepared for the quantification of biochemical parameters. Laboratory tests were performed at the hospital's central laboratory, using standard techniques.

To determine total cholesterol, high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), triglycerides and blood glucose, blood samples were collected in dry tubes, centrifuged for 30 min at 3500 g and quantified by enzymatic techniques using an AU5400 automated analyzer (Beckman Coulter). Biochemical risk markers – lipoprotein(a), apolipoprotein B and high-sensitivity C-reactive protein (hs-CRP) – were quantified with the same analyzer using immunoturbidimetry. Fibrinogen was measured in blood collected into a sodium citrate tube with the subject in a fasting state, using an ACL TOP 700 analyzer (Beckman Coulter).

Genetic analysis of the C825T polymorphism of the *GNB3* gene

Genomic DNA was extracted from peripheral blood leukocytes using the salting-out method. Genotyping was performed using oligonucleotide probes labeled with allele-specific fluorescence in an assay combining standard polymerase chain reaction (PCR) and TaqMan techniques (Applied Biosystems). Primers and probes for the C825T polymorphism of the *GNB3* gene (rs5443) were those provided by the manufacturer (TaqMan SNP Genotyping Assays, Applied Biosystems). Oligonucleotides were synthesized and FAMTM and VIC[®] fluorescent labels were coupled to the 5' ends of the probes for allelic discrimination. Two-step PCR, consisting of 40 denaturation cycles at 92 °C for 15 s and primer annealing and extension at 60 °C for 1 min, was carried out in a 7300 Real-Time PCR System (Applied Biosystems). 7300 System SDS Software (Applied Biosystems) was used to determine genotypes, without using prior knowledge of individual clinical data (Figure 1).

Statistical analysis

Categorical variables are presented as frequency and percentage using the chi-square test. Continuous variables are expressed as mean ± standard deviation or median (minimum-maximum). The Student's t test or the non-parametric Mann-Whitney test was used to compare continuous variables, as appropriate.

Hardy-Weinberg equilibrium was analyzed for each of the genes using the chi-square test and a p-value less than 5% was considered statistically significant. The Bonferroni correction was applied. Multivariate logistic regression analysis using the forward Wald method was applied to identify variables associated with the onset of hypertension.

The strength of associations was determined by odds ratios (OR) and respective 95% confidence intervals (CIs).

The statistical analysis was performed with IBM SPSS version 19.0 and a p value <0.05 was taken to indicate statistical significance.

Results

Characteristics of the study population

The baseline characteristics of the study population are presented in Table 1. The sample consisted of 1641 individuals with hypertension (mean age 50.8±8.1 years; 51.8% male) and 793 controls without hypertension (mean age 50.3±8.2 years; 47.9% male). There were no significant differences between the groups in terms of gender or age.

Significant differences between groups were seen in alcohol consumption (p=0.008), BMI and obesity (p<0.0001), waist circumference (p<0.0001) and waist/hip ratio (p<0.0001), in all of which mean values were higher in the hypertensive group than in the control group. Likewise, diabetes, dyslipidemia, and high SBP, DBP, heart rate and PWV were significantly more frequent in the hypertensive group than in the normotensive group (p<0.0001). In the hypertensive group, there were more individuals with a personal (p=0.002) and family (p<0.0001) history of cardiovascular events, as well as a family history of hypertension (p<0.0001). There was a significantly larger proportion of smokers in the control group (p<0.0001), except in those who smoked >20 cigarettes/day.

Regarding biochemical variables (Table 2), there were significant differences between the groups in hemoglobin, leukocytes, fibrinogen, HDL cholesterol, triglycerides, apolipoprotein B, fasting blood glucose and hs-CRP (all p<0.0001), with higher mean values in the hypertensive group than in controls. There were no significant differences in relation to total cholesterol, LDL cholesterol, lipoprotein(a) and platelets (p>0.05).

Allele and genotype frequencies of the C825T polymorphism of the *GNB3* gene and risk of hypertension

The allele frequencies of the C825T polymorphism of the *GNB3* gene are presented in Table 3. The frequencies of the C and T alleles in the overall population were 60.5% and 39.5%, respectively. The proportions were similar in both cases and controls, with OR 1.106 and without statistical significance (p=0.158) (Table 3).

The frequency of each genotype in both groups was in Hardy-Weinberg equilibrium (Table 4). In the additive model, the most frequent genotype was heterozygous CT, with a frequency of 50.7% in the hypertensive group and 44.3% in the controls compared to the reference homozygous CC genotype. This genotype carried a 1.336-fold greater risk of hypertension (p=0.008) than the mutated homozygote TT (OR 1.107; p=0.494).

In the dominant model (Table 5), subjects with the CT or TT genotype were at greater risk of developing hypertension (OR 1.275; 95% CI 1.042-1.559) than those with the reference genotype (CC) (p=0.018).

Logistic regression analysis

In the logistic regression analysis (Table 6), the variables that remained in the model were diabetes, obesity, alcohol




Figure 1 Genotype determination of the C825T polymorphism of the *GNB3* gene using the 7300 Real-Time PCR System (Applied Biosystems).

Table 1 Demographic and clinical characteristics of the study population.

Variable	Total (n=1641)	Hypertensives (n=848)	Controls (n=793)	p
Age, years	50.6±8.1	50.8±8.1	50.3±8.2	0.212
Male, n (%)	819 (49.9)	439 (51.8)	380 (47.9)	0.119
Sedentary lifestyle, n (%)	887 (54.1)	476 (56.1)	411 (51.8)	0.080
Drinkers, n (%)	627 (38.2)	350 (41.3)	277 (34.9)	0.008
Excessive drinkers ^a , n (%)	536 (32.8)	291 (34.3)	245 (31.2)	0.176
Smokers, n (%)	375 (22.9)	164 (19.5)	211 (26.6)	<0.0001
1-10 ^b , n (%)	142 (37.9)	59 (36)	83 (39.3)	0.002
10-20 ^b , n (%)	118 (31.5)	40 (24.4)	78 (37)	
>20 ^b , n (%)	115 (30.7)	65 (39.6)	50 (23.7)	
BMI, kg/m ²	27.7±4.9	29.2±5.2	26.2±4	<0.0001
Normal ^c , n (%)	466 (28.4)	143 (16.9)	323 (40.7)	<0.0001
Overweight ^d , n (%)	734 (44.7)	379 (44.7)	355 (44.8)	
Obese ^e , n (%)	441 (26.9)	326 (38.4)	115 (14.5)	
High WC ^f , n (%)	1217 (74.2)	710 (83.7)	507 (63.9)	<0.0001
High WHR ^g , n (%)	1461 (89)	788 (92.9)	673 (84.9)	<0.0001
Diabetes, n (%)	169 (10.3)	134 (15.8)	35 (4.4)	<0.0001
Dyslipidemia, n (%)	815 (49.7)	470 (55.4)	345 (43.5)	<0.0001
SBP, mmHg	134.4±20.4	147.2±18.9	120.7±10.9	<0.0001
DBP, mmHg	84.5±12.1	91.1±11.7	77.4±7.7	<0.0001
Heart rate, bpm	72.1±11.8	73.1±12.2	71±11.2	<0.0001
PWV, m/s	8±1.5	8.4±1.5	7.7±1.3	<0.0001
PWV >10 m/s, n (%)	160 (9.8)	117 (13.8)	43 (5.4)	<0.0001
CV events, n (%)	41 (2.5)	31 (3.7)	10 (1.3)	0.002
Family CV history, n (%)	539 (32.9)	327 (38.6)	212 (26.7)	<0.0001
Family hypertension history, n (%)	609 (37.1)	405 (47.8)	204 (25.7)	<0.0001

BMI: body mass index; CV: cardiovascular; DBP: diastolic blood pressure; PWV: pulse wave velocity; SBP: systolic blood pressure; WC: waist circumference; WHR: waist/hip ratio.

^a >70 g/week in men and >50 g/week in women.

^b No. of cigarettes smoked per day.

^c <25 kg/m².

^d 25-29.9 kg/m².

^e ≥30 kg/m².

^f >94 cm in men and >80 cm in women.

^g >0.9 in men and >0.8 in women.

Table 2 Biochemical characteristics of the study population.

Variable	Total (n=1641)	Hypertensives (n=848)	Controls (n=793)	p
Hemoglobin, g/dl	14.3 (9.6-18.2)	14.4 (9.6-18.2)	14.2 (10.1-17.6)	<0.0001
Platelets, 10 ³ /μl	229.0 (23-664)	233.0 (23-664)	227.0 (65-544)	0.206
Leukocytes, 10 ³ /μl	6.4 (2.1-18)	6.6 (2.9-18)	6.2 (2.1-16.6)	<0.0001
Fibrinogen, mg/dl	362 (179.2-874)	371.0 (179.2-874)	355.3 (224-688)	<0.0001
Blood glucose, mg/dl	95 (66-364)	98.0 (70-360)	93.0 (66-364)	<0.0001
Lipoprotein(a), mg/dl	16.3 (0.6-236)	16.6 (0.6-236)	16.2 (0.8-198.2)	0.730
Total cholesterol, mg/dl	207 (107-370)	208.5 (115-344)	206 (107-370)	0.193
HDL-C, mg/dl	48 (17.2-111.7)	46.9 (17.2-103)	49 (20.8-111.7)	<0.0001
LDL-C, mg/dl	131 (37.7-269)	131 (37.7-269)	131 (42-260)	0.944
Triglycerides, mg/dl	110 (21-1098)	119.0 (29-1098)	99.0 (21-688)	<0.0001
Apolipoprotein B, mg/dl	104 (3.9-232)	106.6 (5.1-205)	99.7 (3.9-232)	<0.0001
hs-CRP, mg/dl	0.2 (0-18.5)	0.2 (0-13.1)	0.2 (0-18.5)	<0.0001

Values are presented as median (minimum-maximum).

HDL-C: high-density lipoprotein cholesterol; hs-CRP: high-sensitivity C-reactive protein; LDL-C: low-density lipoprotein cholesterol.

Table 3 Allele frequencies of the C825T polymorphism of the *GNB3* gene.

Allele	Total	%	Cases	%	Controls	%	OR (95% CI)	p
C	1985	60.5	1006	59.3	979	61.7	1.106 (0.962-1.273)	0.158
T	1297	39.5	690	40.7	607	38.3		
Total	3282	100	1696	100	1586	100		

CI: confidence interval; OR: odds ratio.

Table 4 Genotypes of the C825T polymorphism of the *GNB3* gene and risk of hypertension.

Genotype	Cases (n=848)	Controls (n=793)	OR (95% CI)	p
CC	288 (34.0%)	314 (39.6%)	Reference	-
CT	430 (50.7%)	351 (44.3%)	1.336 (1.079-1.653)	0.008
TT	130 (15.3%)	128 (16.1%)	1.107 (0.827-1.482)	0.494

CI: confidence interval; OR: odds ratio.

Table 5 Genotypes of the C825T polymorphism of the *GNB3* gene and risk of hypertension (dominant model).

Genotype (dominant model)	Cases (n=848)	Controls (n=793)	OR (95% CI)	p
CC	288 (34.0%)	314 (39.6%)	Reference	-
CT + TT	560 (66.0%)	479 (40.4%)	1.275 (1.042-1.559)	0.018

CI: confidence interval; OR: odds ratio.

Table 6 Logistic regression analysis of the C825T polymorphism of the *GNB3* gene and confounding variables.

Variable	Beta	SE	Wald	df	OR (95% CI)	p
Diabetes	1.223	0.205	35.711	1	3.397 (2.275-5.074)	<0.0001
Obesity	1.185	0.126	88.686	1	3.270 (2.555-4.184)	<0.0001
Alcohol consumption	0.243	0.109	5.010	1	1.276 (1.031-1.579)	0.025
Smoking	-0.464	0.127	13.359	1	0.629 (0.490-0.806)	<0.0001
Polymorphism (CT+TT)	0.236	0.108	4.755	1	1.267 (1.024-1.566)	0.029
Constant	-0.477	0.102	21.881	1	0.621	<0.0001

Forward Wald method in SPSS version 19.0. Sedentary lifestyle, gender and age left the equation. Statistical significance for p<0.05.

CI: confidence interval; df: degrees of freedom; OR: odds ratio; SE: standard error.

consumption and the CT and TT genotypes of the C825T polymorphism of the *GNB3* gene, which were significantly and independently associated with the occurrence of hypertension.

The model showed smoking to be a protective factor, while a sedentary lifestyle left the model and did not influence the occurrence of hypertension.

Discussion

In the present study, we investigated the association between the 825T variant of the *GNB3* G protein gene and hypertension in a Portuguese population from the Madeira archipelago. Subjects with the dominant genetic model had an increased risk of developing hypertension (OR 1.275; 95% CI 1.042-1.559; $p=0.018$), even after multivariate analysis and correction for confounding factors (obesity, diabetes, sedentary lifestyle, alcohol consumption and smoking). This demonstrates that in this population G protein variants are independently associated with hypertension, constituting an additional risk to conventional risk factors.

The frequency of the T allele was 39.5% in our study, which is higher than that reported in other Caucasian populations (37.7%), but lower than that described for Asians (48.4%) and African-Americans (76.3%).¹⁹

Our results, which are innovative in the Portuguese population, are in line with those of other studies. Stiffer et al.,¹⁴ in a study of 427 normotensive and 426 hypertensive German subjects, were the first to demonstrate a significant association between the C825T polymorphism of the *GNB3* gene coding for pertussis toxin-sensitive G proteins and hypertension (OR 1.79; 95% CI 1.05-3.05; $p=0.03$). Subsequently, other studies in different countries and ethnic groups also found associations between the C825T polymorphism and hypertension^{14,15,17-19} and variations in blood pressure.³²⁻³⁴

In a study of Caucasian subjects, Benjafield et al.¹⁵ concluded that the 825T allele was associated with hypertension (OR 2.3; 95% CI 1.5-3.7),¹⁵ while a study in an Egyptian population also found an association between this variant and hypertension (OR 1.4; 95% CI 0.9-2.1) in the dominant genetic model, which was significantly stronger in the recessive model (OR 2.3; 95% CI 1.5-3.7).²⁰

Zheng et al.¹⁹, in a meta-analysis involving 36 802 individuals, confirmed a significant association between the C825T polymorphism and overall risk of hypertension in Caucasian and Chinese subjects, although this association was not found in other Asian populations.¹⁹

By contrast, a meta-analysis by Guo et al. suggested that although the 825T allele showed a marginal association with hypertension risk, there was no association with hypertension in Asians or Caucasians.²⁶

Various pathophysiological mechanisms may explain the role of the C825T (rs5443) genetic variant in the development of hypertension.

Siffert et al.¹⁴ reported that the splice variant *GNB3-s* is a functional protein that increases G protein activity in a reconstituted system and leads to enhanced intracellular signaling.

Elevated blood pressure is due to increased sensitivity to vasopressor hormones, which use *GNB3* for signal transmission.³⁵ This has been demonstrated in vivo by the greater vascular reactivity shown by individuals with

the 825T variant following stimulation of coronary alpha2 adrenoceptors.³⁶

Meirhaeghe et al.¹⁶ assessed coronary artery vasomotility after intravenous injection of methylethylergonovine maleate, a vasoconstrictor, and concluded that subjects bearing at least one T allele of the C825T polymorphism had greater susceptibility to vasoconstriction than those with the wild-type CC genotype. The *GNB3-s* subunit is associated with increased G protein activity and therefore may increase the activity of alpha-1A adrenoceptors coupled to these proteins. This would explain the greater vasoconstriction observed in response to a vasoconstrictor in individuals with the 825T allele.

Another hypothesis, put forward by Siffert et al.,¹⁴ is that the onset of hypertension in carriers of the T allele may result not from increased vasoconstriction due to vasoconstrictor hormones such as noradrenaline and angiotensin II activating heterotrimeric non-pertussis toxin-sensitive G proteins, but from vascular smooth muscle cell proliferation leading gradually to vascular hypertrophy.

We performed logistic regression analysis to assess confounding variables such as diabetes, alcohol consumption, smoking and the dominant model of the C825T polymorphism, in order to determine which variables were significantly and independently related to the occurrence of hypertension. Obesity and diabetes conferred a significant risk (OR 3.270; 95% CI 2.555-4.184; $p<0.0001$ and OR 3.397; 95% CI 2.275-5.074; $p<0.0001$, respectively). The C825T polymorphism also remained in the model, which revealed that it was significantly and independently associated with the occurrence of hypertension (OR 1.276; 95% CI 1.024-1.566; $p=0.029$).

In our study population there was a larger proportion of smokers (except those who smoked >20 cigarettes/day) in the control group than in the hypertensive group. Multivariate analysis revealed that smoking appeared to be a protective factor for the occurrence of hypertension. This is because smokers were defined as those who smoked at enrollment in the study, so these individuals were active rather than former smokers. Hypertensive individuals were already under medical monitoring for cardiovascular risk factors at enrollment, which would have led some of them to quit smoking. However, our results indicate that the heaviest smokers found it most difficult to quit.

Obesity is also known to be an important risk factor for the development of hypertension. Cross-sectional studies have shown that obesity is associated with higher blood pressure and weight gain throughout life and is an important predictor of hypertension.³⁷⁻⁴⁰ Studies in different ethnic groups have shown that the T allele of the C825T polymorphism is associated with obesity^{41,42} and can influence blood pressure variations.^{43,44}

Future studies that correlate abnormalities in complex cellular transduction mechanisms with other pathological processes may lead to important discoveries in the pharmacological treatment of hypertension and many other diseases.

Strong points of the study

This is the first case-control study conducted in the genetically homogeneous⁴⁵⁻⁴⁷ and relatively isolated Portuguese

population of the archipelago of Madeira to investigate the association between the C825T polymorphism of the *GNB3* gene and propensity for hypertension.

The population's characteristics are advantageous for gene mapping in rare disorders, and furthermore, according to several researchers, the study of a culturally and genetically isolated population with a common way of life, diet and natural environment can help minimize the influence of environment factors on variation in disease patterns.⁴⁸

Conclusion

In this study, the C825T polymorphism of the *GNB3* gene was significantly and independently associated with the occurrence of hypertension in a Portuguese population (OR 1.275; 95% CI 1.042-1.559; $p=0.018$) in the dominant model, even after logistic regression analysis which included other variables that could also have been associated with the onset of hypertension.

Identifying candidate genes and understanding their function will help establish a specific molecular basis for hypertension. Further studies in different populations and with larger sample sizes are needed to determine whether the C825T polymorphism plays a significant role in the etiology of hypertension.

Funding

This study was funded by the Operational Program for the Enhancement of Economic Potential and Territorial Cohesion of the Autonomous Region of Madeira (INTERVIR+).

Conflicts of interest

The authors have no conflicts of interest to declare.

References

- Kearney PM, Whelton M, Reynolds K, et al. Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365:217–23.
- National Heart, Lung and Blood Institute. *Fact Book Fiscal Year*. Bethesda, MD: NHLBI; 2006.
- Lewington S, Clarke P, Qizilbash N, et al., Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360:1903–13.
- Polónia J, Martins L, Pinto F, et al. Prevalence, awareness, treatment and control of hypertension and salt intake in Portugal: changes over a decade. The PHYSA study. *J Hypertens*. 2014;32:1211–21.
- Gong M, Hubner N. Molecular genetics of human hypertension. *Clin Sci*. 2006;110:315–26.
- Lifton RP. Molecular genetics of human blood pressure variation. *Science*. 1996;272:676–80.
- Neves SR, Ram PT, Iyengar R. G protein pathways. *Science*. 2002;296:1636–9.
- Emala CW, Schwindinger WF, Wand GS, et al. Signal-transducing G proteins: basic and clinical implications. *Prog Nucleic Acid Res Mol Biol*. 1994;47:81–111.
- Farfel Z, Bourne HR, Iiri T. The expanding spectrum of G protein diseases. *N Engl J Med*. 1999;340:1012–20.
- Milligan G, Kostenis E. Heterotrimeric G-proteins: a short history. *Br J Pharmacol*. 2006;147:546–55.
- Bezerra MGT, Latronico AC, Fragoso MCBV. Tumores endócrinos associados às mutações das proteínas Gs alfa e Gi alfa2. *Arq Bras Endocrinol Metab*. 2005;49:784–90.
- McCuden CR, Hains MD, Kimple RJ, et al. G-protein signaling: back to the future. *Cell Mol Life Sci*. 2005;62:551–7.
- Hauache OM. Receptores acoplados à proteína G: implicações para a fisiologia e doenças endócrinas. *Arq Bras Endocrinol Metab*. 2001;45:228–39.
- Siffert W, Roszkopf D, Siffert G, et al. Association of a human G-protein $\beta 3$ subunit variant with hypertension. *Nat Genet*. 1998;18:45–8.
- Benjafield AV, Jeyasingam CL, Nyholt DR, et al. G-protein $\beta 3$ subunit gene (*GNB3*) variant in causation of essential hypertension. *Hypertension*. 1998;32:1094–7.
- Meirhaeghe A, Bauters C, Helbecqve N, et al. The human G-protein beta3 subunit C825T polymorphism is associated with coronary artery vasoconstriction. *Eur Heart J*. 2001;22:845–8.
- Siffert W. Effects of the G protein beta 3-subunit gene C825T polymorphism: should hypotheses regarding the molecular mechanisms underlying enhanced G protein activation be revised? Focus on "A Splice Variant of the G Protein Beta 3-Subunit Implicated in Disease States Does Not Modulate Ion Channels". *Physiol Genomics*. 2003;13:81–4.
- Li M, Zhang B, Li C, et al. G-protein beta 3 subunit polymorphisms and essential hypertension: a case-control association study in northern Han Chinese. *J Geriatr Cardiol*. 2015;12:127–34.
- Zheng H, Xu H, Cui B, et al. Association between polymorphism of the G-protein $\beta 3$ subunit C825T and essential hypertension: an updated meta-analysis involving 36,802 subjects. *Biol Res*. 2013;46:265–73.
- El Din Hemimi NS, Mansour AA, Abdelsalam MM. Prediction of the risk for essential hypertension among carriers of C825T genetic polymorphism of G protein $\beta 3$ (*GNB3*) Gene. *Biomark Insights*. 2016;11:69–75.
- Ko KD, Kim KK, Suh HS, et al. Associations between the *GNB3* C825T polymorphism and obesity-related metabolic risk factors in Korean obese women. *J Endocrinol Invest*. 2014;37:1117–20.
- Fang L, Zhou C, Bai S, et al. The C825T polymorphism of the G-protein $\beta 3$ gene as a risk factor for depression: a meta-analysis. *PLoS One*. 2015;10:e0132274.
- Semplicini A, Grandi T, Sandonà C, et al. G-protein $\beta 3$ -subunit gene C825T polymorphism and cardiovascular risk: an updated review. *High Blood Press Cardiovasc Prev*. 2015;22:225–32.
- Brand E, Herrmann SM, Nicaud V, et al. The 825C/T polymorphism of the G-protein subunit beta3 is not related to hypertension. *Hypertension*. 1999;33:1175–8.
- Kato N, Sugiyama T, Morna H, et al. G protein $\beta 3$ subunit variant and essential hypertension in Japanese. *Hypertension*. 1998;32:935–8.
- Guo L, Zhang LL, Zheng B, et al. The C825T polymorphism of the G-protein $\beta 3$ subunit gene and its association with hypertension and stroke: an updated meta-analysis. *PLOS ONE*. 2013;8.
- European Society of Hypertension-European Society of Cardiology Guidelines Committee. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. *J Hypertens*. 2003;21:1011–53.
- National Institute of Health, National Heart, Lung, and Blood Institute North American Association for the Study of Obesity. *The Practical Guide: Identification, Evaluation, and Treatment of Overweight and Obesity in Adults*. NHLBI Obesity Education Initiative, NIH Publication Number: 00-4084; 2000.

29. Authors/Task Force Members: Piepoli MF, Hoes AW, Agewall S, et al. 2016 European Guidelines on Cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts) Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J*. 2016;37:2315–81.
30. Asmar R, Benetos A, Topouchian J, et al. Assessment of arterial distensibility by automatic pulse wave velocity measurement: validation and clinical application studies. *Hypertension*. 1995;26:485–90.
31. Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*. 2003;26:3160–7.
32. Hegele RA, Harris SB, Hanley AJG, et al. G protein $\beta 3$ subunit gene variant and blood pressure variation in Canadian Oji-Cree. *Hypertension*. 1998;32:688–92.
33. Schunkert H, Hense HW, Doring A, et al. Association between a polymorphism in the G protein $\beta 3$ subunit gene and lower renin and elevated diastolic blood pressure levels. *Hypertension*. 1998;32:510–3.
34. Hayakawa T, Takamura T, Abe T, et al. Association of the C825T polymorphism of the G-protein $\beta 3$ subunit gene with hypertension, obesity, hyperlipidemia, insulin resistance, diabetes, diabetic complications, and diabetic therapies among Japanese. *Metabolism*. 2007;56:44–8.
35. Schiffrin EL. Intracellular signal transduction for vasoactive peptides in hypertension. *Can J Physiol Pharmacol*. 1994;72:954–62.
36. Baumgart D, Naber C, Haude M, et al. G protein $\beta 3$ subunit 825T allele and enhanced coronary vasoconstriction on $\alpha 2$ -adrenoceptor activation. *Circ Res*. 1999;85:965–9.
37. Kannel WB, Brand N, Skinner JJ Jr, et al. The relation of adiposity to blood pressure and development of hypertension. The Framingham study. *Ann Intern Med*. 1967;67:48–59.
38. Johnson AL, Cornoni JC, Cassel JC, et al. Influence of race, sex and weight on blood pressure behavior in young adults. *Am J Cardiol*. 1975;35:523–30.
39. Mikhail N, Golub MS, Tuck ML. Obesity and hypertension. *Prog Cardiovasc Dis*. 1999;42:39–58.
40. Frohlich ED, Messerli FH, Reisin E, et al. The problem of obesity and hypertension. *Hypertension*. 1983;5:71–8.
41. Siffert W, Forster P, Jöckel K-H, et al. Worldwide ethnic distribution of the G protein $\beta 3$ subunit 825T allele and its association with obesity in Caucasians, Chinese, and black African individuals. *J Am Soc Nephrol*. 1999;10:1921–30.
42. Siffert W, Naber C, Walla M, et al. G protein $\beta 3$ subunit 825T allele and its potential association with obesity in hypertensive individuals. *J Hypertens*. 1999;17:1095–8.
43. Iiri T, Farfel Z, Bourne HR. G-protein diseases furnish a model for the turn-on switch. *Nature*. 1998;394:35–8.
44. Hegele RA, Anderson C, Young TK, et al. G-protein $\beta 3$ subunit gene splice variant and body fat distribution in Nunavut Inuit. *Genome Res*. 1999;9:972–7.
45. Gonçalves R, Freitas A, Branco M, et al. Y-chromosome lineages from Portugal, Madeira and Açores record elements of Sephardim and Berber ancestry. *Ann Hum Genet*. 2005;69:443–54.
46. Brehm A, Pereira L, Kivisild T, et al. Mitochondrial portraits of the Madeira and Açores archipelagos witness different genetic pools of its settlers. *Hum Genet*. 2003;114:77–86.
47. Kristiansson K, Naukkarinen J, Peltonen L. Isolated populations and complex disease gene identification. *Genome Biol*. 2008;9:109.
48. Jorde LP, Wooding SP. Genetic variation, classification and race. *Nat Genet*. 2004;36 Suppl.:528–33.