

## IMAGE IN CARDIOLOGY

**Severe left ventricular outflow tract obstruction****Obstrução grave do trato de saída do ventrículo esquerdo**

Ana Rita Araújo Silva\*, Inês Carmo Mendes, Susana Cordeiro, Rui Anjos




Serviço de Cardiologia Pediátrica, Hospital de Santa Cruz – Centro Hospitalar de Lisboa Ocidental, Lisboa, Portugal


Received 14 August 2015; accepted 24 August 2015

Available online 31 May 2016

A male neonate was born at term. Obstetric ultrasound at 36 weeks showed two hyperechogenic masses, one located in the mid interventricular septum occupying the total thickness of the septum and the other attached to the anterior leaflet of the mitral valve. The baby was born by normal delivery, with a birth weight of 3280 g and an Apgar score of 10 at first minute and 10 at fifth minute. On physical examination, he had a grade III/VI systolic heart murmur best heard at the left sternal edge. An echocardiogram on day one revealed multiple masses with features of rhabdomyomas, located at the apex of the right ventricle, tricuspid and mitral valves, interventricular septum, left ventricular lateral wall and outflow tract. The left ventricular outflow tract mass (15 mm×9 mm) caused a peak gradient of 92 mmHg and a mean gradient of 49 mmHg (Figure 1a and b). The newborn underwent serial clinical and echocardiographic assessment. He remained asymptomatic and the masses underwent progressive reduction. At one month the left ventricular outflow tract mass measured 14 mm×8 mm and at latest follow-up at six months measured 12 mm×8 mm (Figure 2a and b). The peak left ventricle-aorta gradient had decreased to 29 mmHg, with a mean of 13 mmHg, at the last follow-up. A diagnosis of tuberous sclerosis was made by genetic testing, which confirmed a mutation in the TSC-2 gene.



**Figure 1** (a) Day one, apical 5-chamber view: a large mass obstructing the left ventricular outflow tract, measuring 15 mm×9 mm; (b) day one, parasternal long-axis view: a large mass almost completely occluding the left ventricular outflow tract. Ao: aorta; LA: left atrium; LV: left ventricle; RA: right atrium; RV: right ventricle.



**Figure 2** (a) Six months, apical 5-chamber view: regression of the mass obstructing the left ventricular outflow tract, now measuring 12 mm×8 mm; (b) six months, parasternal long-axis view: clear regression of the mass. Ao: aorta; LA: left atrium; LV: left ventricle; RA: right atrium; RV: right ventricle.

\* Corresponding author.

E-mail address: [ana.araujo.silva@hotmail.com](mailto:ana.araujo.silva@hotmail.com)  
(A.R. Araújo Silva).

A small reduction in the volume of obstructive masses in infants leads to a significant decrease in gradients. Sometimes it is better to wait!

### **Ethical disclosures**

**Protection of human and animal subjects.** The authors declare that no experiments were performed on humans or animals for this study.

**Confidentiality of data.** The authors declare that no patient data appear in this article.

**Right to privacy and informed consent.** The authors declare that no patient data appear in this article.

### **Conflicts of interest**

The authors have no conflicts of interest to declare.